

Exercise 11.1

1. If a line makes angles 90° , 135° , 45° with the x , y and z -axes respectively, find its direction cosines.

Sol. We know that direction cosines of a line making angles α , β , γ with the x , y and z -axes respectively are $\cos \alpha$, $\cos \beta$, $\cos \gamma$.

Here $\alpha = 90^\circ$, $\beta = 135^\circ$ and $\gamma = 45^\circ$.

Therefore, direction cosines of the required line are $\cos 90^\circ$,

$\cos 135^\circ$ and $\cos 45^\circ = 0$, $\frac{-1}{\sqrt{2}}$ and $\frac{1}{\sqrt{2}}$.

$$\left[\begin{array}{l} \cos 135^\circ = \cos (180^\circ - 45^\circ) = -\cos 45^\circ = \frac{-1}{\sqrt{2}} \\ \text{(II)} \end{array} \right]$$

Result. $\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma = 1$.

2. Find the direction cosines of a line which makes equal angles with the co-ordinate axes.

Sol. Let a line make equal angles α, α, α with the co-ordinate axes.

\therefore Direction cosines of the line are $\cos \alpha, \cos \alpha, \cos \alpha \dots (i)$

$\therefore \cos^2 \alpha + \cos^2 \alpha + \cos^2 \alpha = 1$ [$\because \cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma = 1$]

$$\Rightarrow 3 \cos^2 \alpha = 1 \quad \Rightarrow \cos^2 \alpha = \frac{1}{3} \quad \Rightarrow \cos \alpha = \pm \sqrt{\frac{1}{3}} = \pm \frac{1}{\sqrt{3}}$$

Putting $\cos \alpha = \pm \frac{1}{\sqrt{3}}$ in (i), direction cosines of the required line making equal angles with the co-ordinate axes are

$$\pm \frac{1}{\sqrt{3}}, \pm \frac{1}{\sqrt{3}}, \pm \frac{1}{\sqrt{3}}.$$

Very Important Remark. Therefore, direction cosines of a line making equal angles with the co-ordinate axes in the positive (i.e.,

first) octant are $\frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}}$.

3. If a line has direction ratios - 18, 12, - 4, then what are its direction cosines?

Sol. We know that if a, b, c are direction ratios of a line, then direction cosines of the line are

$$\frac{a}{\sqrt{a^2 + b^2 + c^2}}, \frac{b}{\sqrt{a^2 + b^2 + c^2}}, \frac{c}{\sqrt{a^2 + b^2 + c^2}} \quad \dots (i)$$

Here, direction ratios of the line are

$$- 18, 12, - 4 = a, b, c$$

$$\therefore \frac{a}{\sqrt{a^2 + b^2 + c^2}} = \frac{-18}{\sqrt{(-18)^2 + (12)^2 + (-4)^2}} = \frac{-18}{\sqrt{324 + 144 + 16}} \\ = \frac{-18}{\sqrt{484}} = \frac{-18}{22} = \frac{-9}{11}$$

Putting these values in (i), direction cosines of the required line are

$$\frac{-18}{22}, \frac{12}{22}, \frac{-4}{22} = \frac{-9}{11}, \frac{6}{11}, \frac{-2}{11}.$$

4. Show that the points (2, 3, 4), (- 1, - 2, 1), (5, 8, 7) are collinear.

Sol. The given points are A(2, 3, 4), B(- 1, - 2, 1) and C(5, 8, 7).

\therefore Direction ratios of the line joining A and B are

$$i.e., \frac{-1-2}{-3}, \frac{-2-3}{-5}, \frac{1-4}{-3} \quad | \quad x_2 - x_1, y_2 - y_1, z_2 - z_1 \quad \dots (i) \\ = a_1, b_1, c_1 \quad (\text{say})$$

Again direction ratios of the line joining B and C are

$$5 - (-1), 8 - (-2), 7 - 1 = 6, 10, 6 \quad \dots (ii) \\ = a_2, b_2, c_2 \quad (\text{say})$$

From (i) and (ii) direction ratios of AB and BC are proportional

$$i.e., \frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2} \quad \left[\frac{-3}{6} = \frac{-5}{10} = \frac{-3}{6} \quad (\text{each} = \frac{-1}{2}) \right]$$

$$a^2 + b^2 = c^2 \quad \left[\begin{array}{l} 6^2 + 10^2 = 2^2 \end{array} \right]$$

Therefore, AB^2 is parallel to BC^2 . But point B is common to both AB and BC. Hence, points A, B, C are collinear.



5. Find the direction cosines of the sides of the triangle whose vertices are (3, 5, -4), (-1, 1, 2) and (-5, -5, -2).


Sol. Direction ratios of line AB are $-1 - 3, 1 - 5, 2 - (-4)$

$$\text{i.e., } -4, -4, 6 \quad | \quad x_2 - x_1, y_2 - y_1, z_2 - z_1$$

$$\begin{aligned} \text{Dividing each by } \sqrt{a^2 + b^2 + c^2} &= \sqrt{(-4)^2 + (-4)^2 + 6^2} \\ &= \sqrt{16 + 16 + 36} = \sqrt{68} = \sqrt{4 \times 17} = 2\sqrt{17}. \end{aligned}$$

direction cosines of line AB are

$$\begin{aligned} &\frac{-4}{2\sqrt{17}}, \frac{-4}{2\sqrt{17}}, \frac{6}{2\sqrt{17}} \\ \text{i.e., } &\frac{-2}{\sqrt{17}}, \frac{-2}{\sqrt{17}}, \frac{3}{\sqrt{17}} \end{aligned}$$



Direction ratios of line BC are

$$-5 - (-1), -5 - 1, -2 - 2 = -4, -6, -4$$

$$\begin{aligned} \text{Dividing each by } \sqrt{(-4)^2 + (-6)^2 + (-4)^2} &= \sqrt{16 + 36 + 16} \\ &= \sqrt{68} = \sqrt{4 \times 17} = 2\sqrt{17} \end{aligned}$$

Direction cosines of line BC are $\frac{-4}{2\sqrt{17}}, \frac{-6}{2\sqrt{17}}, \frac{-4}{2\sqrt{17}}$

$$\text{i.e., } \frac{-2}{\sqrt{17}}, \frac{-3}{\sqrt{17}}, \frac{-2}{\sqrt{17}}$$

Direction ratios of line CA are

$$3 - (-5), 5 - (-5), -4 - (-2) = 8, 10, -2$$

$$\begin{aligned} \text{Dividing each by } \sqrt{(8)^2 + (10)^2 + (-2)^2} &= \sqrt{64 + 100 + 4} \\ &= \sqrt{168} = \sqrt{4 \times 42} = 2\sqrt{42}. \end{aligned}$$

Direction ratios of line CA are

$$\frac{8}{2\sqrt{42}}, \frac{10}{2\sqrt{42}}, \frac{-2}{2\sqrt{42}} = \frac{4}{\sqrt{42}}, \frac{5}{\sqrt{42}}, \frac{-1}{\sqrt{42}}$$

Note. If l, m, n are direction cosines of a line, then $-l, -m, -n$ are also direction cosines of the same line.

Exercise 11.2

1. Show that the three lines with direction cosines

$$\frac{12}{13}, \frac{-3}{13}, \frac{-4}{13}; \frac{4}{13}, \frac{12}{13}, \frac{3}{13}; \frac{3}{13}, \frac{-4}{13}, \frac{12}{13}$$

are mutually perpendicular.

Sol. Given: Direction cosines of three lines are

$$\frac{12}{13}, \frac{-3}{13}, \frac{-4}{13}; = l_1, m_1, n_1, \quad \frac{4}{13}, \frac{12}{13}, \frac{3}{13}; = l_2, m_2, n_2$$

and $\frac{3}{13}, \frac{-4}{13}, \frac{12}{13} = l_3, m_3, n_3$

For first two lines;

$$l_1 l_2 + m_1 m_2 + n_1 n_2 = \frac{12}{13} \left(\frac{4}{13} \right) + \frac{(-3)}{13} \left(\frac{12}{13} \right) + \frac{(-4)}{13} \left(\frac{3}{13} \right)$$

$$= \frac{48}{169} - \frac{36}{169} - \frac{12}{169} = \frac{48-36-12}{169} = \frac{0}{169} = 0$$

∴ The first two lines are perpendicular to each other.

For second and third line,

$$l_2 l_3 + m_2 m_3 + n_2 n_3 = \frac{4}{13} \left(\frac{3}{13} \right) + \frac{12}{13} \left(\frac{-4}{13} \right) + \frac{3}{13} \left(\frac{12}{13} \right) = \frac{12}{169} - \frac{48}{169} + \frac{36}{169}$$

$$= \frac{12-48+36}{169} = \frac{0}{169} = 0$$

∴ Second and third lines are perpendicular to each other.

For first and third line,

$$l_1 l_3 + m_1 m_3 + n_1 n_3 = \frac{12}{13} \left(\frac{3}{13} \right) + \frac{(-3)}{13} \left(\frac{-4}{13} \right) + \frac{(-4)}{13} \left(\frac{12}{13} \right) = \frac{36}{169} + \frac{12}{169} - \frac{48}{169}$$

$$= \frac{36+12-48}{169} = \frac{0}{169} = 0$$

∴ First and third line are also perpendicular to each other.

∴ The three given lines are mutually perpendicular.

2. **Show that the line through the points (1, -1, 2), (3, 4, -2) is perpendicular to the line through the points (0, 3, 2) and (3, 5, 6).**

Sol. We know that direction ratios of the line joining the points

A(1, -1, 2) and B(3, 4, -2) are $x_2 - x_1, y_2 - y_1, z_2 - z_1$
i.e., $3 - 1, 4 - (-1), -2 - 2 = 2, 5, -4 = a_1, b_1, c_1$ (say)
 Again, direction ratios of the line joining the points

C(0, 3, 2) and D(3, 5, 6) are $x_2 - x_1, y_2 - y_1, z_2 - z_1$
i.e., $3 - 0, 5 - 3, 6 - 2 = 3, 2, 4 = a_2, b_2, c_2$ (say)

For these lines AB and CD,

$$a_1 a_2 + b_1 b_2 + c_1 c_2 = 2(3) + 5(2) + (-4)(4)$$

$$= 6 + 10 - 16 = 0$$

∴ Given line AB is perpendicular to given line CD.

3. **Show that the line through the points (4, 7, 8), (2, 3, 4) is parallel to the line through the points (-1, -2, 1), (1, 2, 5).**

Sol. We know that direction ratios of the line joining the points

A(4, 7, 8) and B(2, 3, 4) are $x_2 - x_1, y_2 - y_1, z_2 - z_1$

i.e., $2 - 4, 3 - 7, 4 - 8$ *i.e.*, $-2, -4, -4 = a_1, b_1, c_1$ (say)
Again, direction ratios of the line joining the points $C(-1, -2, 1)$
and $D(1, 2, 5)$ are $1 - (-1), 2 - (-2), 5 - 1 = 2, 4, 4 = a_2, b_2, c_2$
(say)

For these lines AB and CD,



$$\left(\text{as } \frac{-2}{1} = \frac{-4}{2} = \frac{-4}{4} (= -1 \text{ each}) \right)$$

$$\underline{a}_1 = \underline{b}_1 = \underline{c}_1$$

$$\therefore \underline{a}_2 \quad \underline{b}_2 \quad \underline{c}_2 \quad \left(\begin{array}{ccc} 2 & 4 & 4 \end{array} \right)$$

Given line AB is parallel to given line CD.


4. Find the equation of the line which passes through the point (1, 2, 3) and is parallel to the vector

$$\hat{i} \quad \hat{j} \quad \hat{k} \quad A(1, 2, 3)$$

$$3\hat{i} + 2\hat{j} - 2\hat{k}$$

Sol. A point on the required line is $A(1, 2, 3) = (x_1, y_1, z_1)$

i.e., Position vector of a point on the required line is



$$\vec{a} = \vec{OA} = (1, 2, 3) = \hat{i} + 2\hat{j} + 3\hat{k}$$

The required line is parallel to the vector $\vec{b} = 3\hat{i} + 2\hat{j} - 2\hat{k}$ (and hence direction ratios of the required line are coefficient of $\hat{i}, \hat{j}, \hat{k}$ in \vec{b} i.e., $3, 2, -2 = a, b, c$)

∴ Vector equation of required line is

$$\vec{r} = \vec{a} + \lambda \vec{b} \quad \text{i.e., } \vec{r} = (\hat{i} + 2\hat{j} + 3\hat{k}) + \lambda(3\hat{i} + 2\hat{j} - 2\hat{k})$$

where λ is a real number.

Remark. Also cartesian equation of the required line in this Q. No. 4 is

$$\frac{x-x_1}{a} = \frac{y-y_1}{b} = \frac{z-z_1}{c} \quad \text{i.e., } \frac{x-1}{3} = \frac{y-2}{2} = \frac{z-3}{-2}$$

5. Find the equation of the line in vector and in cartesian form that passes through the point with position vector

$$2\hat{i} - \hat{j} + 4\hat{k} \quad \text{and is in the direction } \hat{i} + 2\hat{j} - \hat{k}$$

Sol. Position vector of a point on the required line is

$$\vec{a} = 2\hat{i} - \hat{j} + 4\hat{k} = (2, -1, 4) = (x_1, y_1, z_1)$$

The required line is in the direction of the vector

$$\vec{b} = \hat{i} + 2\hat{j} - \hat{k}$$

⇒ direction ratios of required line are coefficients of \hat{i} , \hat{j} , \hat{k}
 →

in b i.e., $1, 2, -1 = a, b, c$

∴ Equation of the required line in vector form is $\vec{r} = \vec{a} + \lambda \vec{b}$
 → ^ ^ ^ ^ ^ ^

i.e., $\vec{r} = (2\hat{i} - \hat{j} + 4\hat{k}) + \lambda(\hat{i} + 2\hat{j} - \hat{k})$

where λ is a real number and equation of line in cartesian form is

$$\frac{z-z_1}{a} = \frac{y-y_1}{b} = \frac{x-x_1}{c} \quad \text{i.e.,} \quad \frac{z-2}{1} = \frac{y+1}{2} = \frac{x-4}{-1}$$




6. Find the cartesian equation of the line which passes through the point $(-2, 4, -5)$ and parallel to the line given by

$$\frac{x+3}{3} = \frac{y-4}{5} = \frac{z+8}{6}$$

Sol. Given: A point on the required line is $(-2, 4, -5) = (x_1, y_1, z_1)$.
Equations of the given line in cartesian form are

$$\frac{z+3}{3} = \frac{y-4}{5} = \frac{x+8}{6}$$



(It is standard form because coefficients of x, y, z are unity each)

\therefore Direction ratios (D.R.'s) of the given line are its denominators 3, 5, 6 and hence d.r.'s of the required parallel line are also 3, 5, 6 = a, b, c .

\therefore Equations of the required line are

$$\frac{z-z_1}{a} = \frac{y-y_1}{b} = \frac{x-x_1}{c} \quad \text{i.e.,} \quad \frac{z-(-2)}{3} = \frac{y-4}{5} = \frac{x-(-5)}{6}$$

$$\text{i.e.,} \quad \frac{z+2}{3} = \frac{y-4}{5} = \frac{x+5}{6}$$

7. The cartesian equation of a line is $\frac{x-5}{3} = \frac{y+4}{7} = \frac{z-6}{2}$.

Write its vector form.

Sol. Given: The cartesian equation of a line is

$$\frac{z-5}{3} = \frac{y+4}{7} = \frac{x-6}{2}$$

$$\text{i.e.} \quad \frac{z-5}{3} = \frac{y-(-4)}{7} = \frac{z-6}{2}$$

comparing the given equation with the standard form

$$\frac{x-x_1}{a} = \frac{y-y_1}{b} = \frac{z-z_1}{c}$$

we have $x_1 = 5, y_1 = -4, z_1 = 6; a = 3, b = 7, c = 2$

Hence the given line passes through the point

$$\vec{a} = (x_1, y_1, z_1) = (5, -4, 6) = 5\hat{i} - 4\hat{j} + 6\hat{k}$$

and is parallel (or collinear) with the vector



$$\vec{b} = a\hat{i} + b\hat{j} + c\hat{k} = 3\hat{i} + 7\hat{j} + 2\hat{k}$$

∴ Vector equation of the given line is $\vec{r} = \vec{a} + \lambda \vec{b}$

$$\text{i.e } \vec{r} = (5\hat{i} - 4\hat{j} + 6\hat{k}) + \lambda (3\hat{i} + 7\hat{j} + 2\hat{k})$$



8. Find the vector and cartesian equations of the line that passes through the origin and (5, -2, 3).

Sol. Vector equation of the line

\vec{a} = Position vector of a point here O (say) on the line

$$= (0, 0, 0) = 0\hat{i} + 0\hat{j} + 0\hat{k} = \vec{0}$$

\vec{b}

= A vector along the line

\vec{b}

= OA = Position vector of point A – Position vector of point O

$$= (5, -2, 3) - (0, 0, 0) = (5, -2, 3) = 5\hat{i} - 2\hat{j} + 3\hat{k}$$

\therefore Vector equation of the line is $\vec{r} = \vec{a} + \lambda \vec{b}$

i.e., $\vec{r} = \vec{0} + \lambda(5\hat{i} - 2\hat{j} + 3\hat{k})$ i.e. $\vec{r} = \lambda(5\hat{i} - 2\hat{j} + 3\hat{k})$.

Cartesian equation of the line

Direction ratios of line OA are 5 - 0, -2 - 0, 3 - 0

i.e., 5, -2, 3

| $x_2 - x_1, y_2 - y_1, z_2 - z_1 = a, b, c$

A point on the line O is $(0, 0, 0) = (x_1, y_1, z_1)$.

\therefore Cartesian equation of the line is

$$\frac{z - z_1}{a} = \frac{y - y_1}{b} = \frac{x - x_1}{c} \quad \text{i.e.,} \quad \frac{z - 0}{5} = \frac{y - 0}{-2} = \frac{x - 0}{3}$$

i.e., $\frac{z}{5} = \frac{y}{-2} = \frac{x}{3}$.

Remark. In the solution of the above question we can also take:

\vec{a} = Position vector of point A = $(5, -2, 3) = 5\hat{i} - 2\hat{j} + 3\hat{k}$

for vector form and point A as $(x_1, y_1, z_1) = (5, -2, 3)$ for Cartesian form.

The equation of line in vector form is $\vec{r} = \vec{a} + \lambda \vec{b}$

$$\vec{r} = 5\hat{i} - 2\hat{j} + 3\hat{k} + \lambda(5\hat{i} - 2\hat{j} + 3\hat{k})$$

i.e., $\vec{r} = 5\hat{i} - 2\hat{j} + 3\hat{k} + \lambda(5\hat{i} - 2\hat{j} + 3\hat{k})$

and equation of line in Cartesian form is $\frac{z-5}{5} = \frac{y+2}{-2} = \frac{x-3}{3}$.

9. Find vector and cartesian equations of the line that passes through the points $(3, -2, -5)$ and $(3, -2, 6)$.

Sol. Vector Equation

Let \vec{a} and \vec{b} be the position vectors of the points $A(3, -2, -5)$ and $B(3, -2, 6)$.

$$\therefore \vec{a} = 3\hat{i} - 2\hat{j} - 5\hat{k} \text{ and } \vec{b} = 3\hat{i} - 2\hat{j} + 6\hat{k}$$



\therefore A vector along the line = \vec{AB} = position vector of point B - position vector of point A

$$= \vec{b} - \vec{a} = 3\hat{i} - 2\hat{j} + 6\hat{k} - 3\hat{i} + 2\hat{j} + 5\hat{k} = 11\hat{k}$$

Vector equation of the line is

$$\vec{r} = \vec{a} + \lambda \vec{AB} \quad \text{i.e.,} \quad \vec{r} = 3\hat{i} - 2\hat{j} - 5\hat{k} + \lambda(11\hat{k})$$

$$\text{i.e.,} \quad \vec{r} = 3\hat{i} - 2\hat{j} - 5\hat{k} + 11\lambda\hat{k}.$$

Note. Another vector equation for the same line is

$$\vec{r} = \vec{b} + \lambda \vec{AB} \quad \text{i.e.,} \quad \vec{r} = 3\hat{i} - 2\hat{j} + 6\hat{k} + 11\lambda\hat{k}.$$

Cartesian Equation

Direction ratios of line AB are 3 - 3, - 2 + 2, 6 + 5

$$\text{i.e.,} \quad 0, 0, 11 \quad | \quad x_2 - x_1, y_2 - y_1, z_2 - z_1$$

\therefore Equations of the line are $\frac{z-z_1}{a} = \frac{y-y_1}{b} = \frac{x-x_1}{c}$

$$\text{i.e.,} \quad \frac{z-3}{0} = \frac{y+2}{0} = \frac{x+5}{11}$$

10. Find the angle between the following pairs of lines:

(i) $\vec{r} = 2\hat{i} - 5\hat{j} + \hat{k} + \lambda(3\hat{i} + 2\hat{j} + 6\hat{k})$ and

$$\vec{r} = 7\hat{i} - 6\hat{k} + \mu(\hat{i} + 2\hat{j} + 2\hat{k})$$

(ii) $\vec{r} = 3\hat{i} + \hat{j} - 2\hat{k} + \lambda(\hat{i} - \hat{j} - 2\hat{k})$ and

$$\vec{r} = 2\hat{i} - \hat{j} - 5\hat{k} + \mu(3\hat{i} - 5\hat{j} - 4\hat{k}).$$

Sol. (i) **Given:** Equation of one line is

$$\vec{r} = 2\hat{i} - 5\hat{j} + \hat{k} + \lambda(3\hat{i} + 2\hat{j} + 6\hat{k})$$

Comparing with $\vec{r} = \vec{a}_1 + \lambda \vec{b}_1$,

$$\vec{a}_1 = 2\hat{i} - 5\hat{j} + \hat{k} \quad \vec{b}_1 = 3\hat{i} + 2\hat{j} + 6\hat{k}$$



$$\vec{r}_1 = 3\hat{i} - 5\hat{j} + \hat{k} \quad \text{and a vector along the line is} \\ \vec{b}_1 = 2\hat{j} + 6\hat{k} \quad \rightarrow \quad \dots(i)$$

(It may be noted that vector \vec{a}_1 is the position vector of a point on the line and not a vector along the line).

Given: Equation of second line is

$$\vec{r} = 7\hat{i} - 6\hat{k} + \mu(\hat{i} + 2\hat{j} + 2\hat{k})$$

Comparing with $\vec{r} = \vec{a}_2 + \mu\vec{b}_2$ we have

$$\vec{a}_2 = 7\hat{i} - 6\hat{k} \quad \text{and a vector along the second line is} \\ \vec{b}_2 = \hat{i} + 2\hat{j} + 2\hat{k}$$

$$\vec{b}_2 = \hat{i} + 2\hat{j} + 2\hat{k} \quad \dots(ii)$$

Let θ be the angle between the two lines.

$$\begin{aligned} \text{We know that } \cos \theta &= \frac{\vec{b}_1 \cdot \vec{b}_2}{|\vec{b}_1| |\vec{b}_2|} \\ &= \frac{3(1) + 2(2) + 6(2)}{\sqrt{9+4+36} \sqrt{1+4+4}} = \frac{3+4+12}{\sqrt{49} \sqrt{9}} \end{aligned}$$

$$\cos \theta = \frac{19}{7(3)} = \frac{19}{21} \quad \therefore \quad \theta = \cos^{-1} \frac{19}{21}$$

(ii) Comparing the equations of the two given lines with

$$\vec{r} = \vec{a}_1 + \lambda \vec{b}_1 \quad \text{and} \quad \vec{r} = \vec{a}_2 + \mu \vec{b}_2 \quad \text{we have}$$

$$\vec{b}_1 = \hat{i} - \hat{j} - 2\hat{k} \quad \text{and} \quad \vec{b}_2 = 3\hat{i} - 5\hat{j} - 4\hat{k}$$

Let θ be the angle between the two lines

$$\begin{aligned} \therefore \cos \theta &= \frac{\vec{b}_1 \cdot \vec{b}_2}{|\vec{b}_1| |\vec{b}_2|} = \frac{(1)(3) + (-1)(-5) + (-2)(-4)}{\sqrt{1+1+4} \sqrt{9+25+16}} \\ &= \frac{3+5+8}{\sqrt{6} \sqrt{50}} \end{aligned}$$

$$= \frac{16}{\sqrt{300}} = \frac{16}{\sqrt{3 \times 100}} = \frac{16}{10\sqrt{3}}$$

$$\text{or } \cos \theta = \frac{8}{5\sqrt{3}} \quad \therefore \quad \theta = \cos^{-1} \frac{8}{5\sqrt{3}}$$

11. Find the angle between the following pairs of lines:

$$(i) \quad \frac{x-2}{2} = \frac{y-1}{5} = \frac{z+3}{-3} \quad \text{and} \quad \frac{x+2}{-1} = \frac{y-4}{8} = \frac{z-5}{4}$$

$$(ii) \quad \frac{x}{2} = \frac{y}{2} = \frac{z}{1} \quad \text{and} \quad \frac{x-5}{4} = \frac{y-2}{1} = \frac{z-3}{8}$$

Sol. (i) Given: Equation of one line is $\frac{z-2}{2} = \frac{y-1}{5} = \frac{x+3}{-3}$

(It is standard form because coefficients of x, y, z are unity each)

\therefore Denominators 2, 5, -3 are direction ratios of this line
i.e., a vector along the line is

$$\vec{b}_1 = (2, 5, -3) = 2\hat{i} + 5\hat{j} - 3\hat{k} \quad \dots(i)$$

Given: Equation of second line is $\frac{z+2}{-1} = \frac{y-4}{8} = \frac{x-5}{4}$

(It is also standard form)

\therefore Denominators -1, 8, 4 are direction ratios of this line
i.e., a vector along the line is



$$\vec{b}_2 = (-1, 8, 4) = -\hat{i} + 8\hat{j} + 4\hat{k} \quad \dots(ii)$$

Let θ be the angle between the two given lines.

$$\begin{aligned} \vec{b}_1 \cdot \vec{b}_2 \\ \rightarrow \rightarrow \\ \text{We know that } \cos \theta = \frac{\vec{b}_1 \cdot \vec{b}_2}{|\vec{b}_1| |\vec{b}_2|} \\ = \frac{2(-1) + 5(8) + (-3)(4)}{\sqrt{4 + 25 + 9} \sqrt{1 + 64 + 16}} \quad (\text{From (i) and (ii)}) \end{aligned}$$

$$\Rightarrow \cos \theta = \frac{-2 + 40 - 12}{\sqrt{38} \sqrt{81} \sqrt{38}} = \frac{26}{9 \sqrt{38}} \Rightarrow \theta = \cos^{-1} \left(\frac{26}{9 \sqrt{38}} \right)$$

(ii) **Given:** Equation of one line is

$$\frac{z}{2} = \frac{y}{2} = \frac{x}{1} \quad (\text{Standard Form})$$

\therefore Denominators 2, 2, 1 are direction ratios of this line i.e., a vector along this line is

$$\vec{b}_1 = (2, 2, 1) = 2\hat{i} + 2\hat{j} + \hat{k} \quad \dots(i)$$

Given: Equation of second line is

$$\frac{z-5}{4} = \frac{y-2}{1} = \frac{x-3}{8} \quad (\text{Standard Form})$$

\therefore Denominators 4, 1, 8 are direction ratios of this line i.e., a vector along this line is

$$\vec{b}_2 = (4, 1, 8) = 4\hat{i} + \hat{j} + 8\hat{k} \quad \dots(ii)$$

Let θ be the angle between the two lines.

$$\begin{aligned} \vec{b}_1 \cdot \vec{b}_2 \\ \rightarrow \rightarrow \\ \text{We know that } \cos \theta = \frac{\vec{b}_1 \cdot \vec{b}_2}{|\vec{b}_1| |\vec{b}_2|} \\ = \frac{2(4) + 2(1) + 1(8)}{\sqrt{4 + 4 + 1} \sqrt{16 + 1 + 64}} = \frac{8 + 2 + 8}{\sqrt{9} \sqrt{81}} = \frac{18}{3 \times 9} = \frac{2}{3} \end{aligned}$$

$$\therefore \theta = \cos^{-1} \frac{2}{3}$$

12. Find the values of p so that the lines $\frac{1-x}{3} = \frac{7y-14}{2p} = \frac{z-3}{2}$ and $\frac{7-7x}{3p} = \frac{y-5}{1} = \frac{6-z}{5}$ are at right angles.

Sol. Let us put the equations of these lines in standard form (*i.e.*, making coeff. of x, y, z unity in each of them)



The first line can be written as

$$-\frac{(z-1)}{3} = \frac{7(y-2)}{2p} = \frac{x-3}{2} \quad \text{or} \quad \frac{z-1}{-3} = \frac{y-2}{\left(\frac{2p}{7}\right)} = \frac{x-3}{2}$$

$$\left(\downarrow 7 \downarrow \right)$$

\therefore direction ratios of this line are $-3, \frac{2p}{7}, 2 = a_1, b_1, c_1$.

And the equation of 2nd line can be written as

$$\frac{-7(z-1)}{3p} = \frac{y-5}{1} = \frac{-(x-6)}{5} \quad \text{or} \quad \frac{z-1}{-\frac{3p}{7}} = \frac{y-5}{1} = \frac{x-6}{-5}$$

\therefore The direction ratios of 2nd line are $\frac{-3p}{7}, 1, -5 = a_2, b_2, c_2$.

\therefore The two lines are perpendicular, therefore

$$\Rightarrow -3 \left(\frac{-3p}{7} \right) + \left(\frac{2p}{7} \right) + c_1 c_2 = 0$$

$$\left(\downarrow 7 \downarrow \right) \quad \left(\downarrow 7 \downarrow \right)$$

$$\Rightarrow \frac{9p}{7} + \frac{2p}{7} - 10 = 0 \quad \Rightarrow \frac{11p}{7} = 10 \quad \Rightarrow p = \frac{70}{11}$$

13. Show that the lines $\frac{x-5}{7} = \frac{y+2}{-5} = \frac{z}{1}$ and $\frac{x}{1} = \frac{y}{2} = \frac{z}{3}$

are perpendicular to each other.

Sol. Given: Equation of one line is

$$\frac{z-5}{7} = \frac{y+2}{-5} = \frac{x}{1} \quad (\text{Standard form})$$

Direction ratios of this line are its denominators $7, -5, 1$

$$= a_1, b_1, c_1 \quad (\Rightarrow \vec{b}_1 = 7\hat{i} - 5\hat{j} + \hat{k})$$

Given: Equation of second line is $\frac{z}{1} = \frac{y}{2} = \frac{x}{3}$ (Standard form)

Direction ratios of this line are its denominators $1, 2, 3$

$$= a_2, b_2, c_2 \quad (\Rightarrow \vec{b}_2 = \hat{i} + 2\hat{j} + 3\hat{k})$$

$$+ 2 \hat{j} + 3 \hat{k}$$

$$\begin{aligned} \hat{b}_1 \cdot \hat{b}_2 &= a_1 a_2 + b_1 b_2 + c_1 c_2 = 7(1) + (-5)(2) + 1(3) \\ &= 7 - 10 + 3 = 0 \end{aligned}$$

\therefore The two given lines are perpendicular to each other.

14. Find the shortest distance between the lines

$$\vec{r} = (\hat{i} + 2\hat{j} + \hat{k}) + \lambda(\hat{i} - \hat{j} + \hat{k}) \text{ and}$$

$$\vec{r} = 2\hat{i} - \hat{j} - \hat{k} + \mu(2\hat{i} + \hat{j} + 2\hat{k}).$$



Sol. Comparing the equations of the given lines with

$$\vec{r} = \vec{a}_1 + \lambda \vec{b}_1 \quad \text{and} \quad \vec{r} = \vec{a}_2 + \mu \vec{b}_2, \quad \text{we have}$$

$$\vec{a}_1 = \hat{i} + 2\hat{j} + \hat{k}, \quad \vec{b}_1 = \hat{i} - \hat{j} + \hat{k}$$

and $\vec{a}_2 = 2\hat{i} - \hat{j} - \hat{k}, \quad \vec{b}_2 = 2\hat{i} + \hat{j} + 2\hat{k}$

We know that the S.D. between the two skew lines is given by

$$d = \frac{|\vec{b}_2 - \vec{a}_1 \cdot (\vec{b}_1 \times \vec{b}_2)|}{|\vec{b}_1 \times \vec{b}_2|} \quad \dots(i)$$

Now $\vec{a}_2 - \vec{a}_1 = (2\hat{i} - \hat{j} - \hat{k}) - (\hat{i} + 2\hat{j} + \hat{k}) = \hat{i} - 3\hat{j} - 2\hat{k}$

$$\vec{b}_1 \times \vec{b}_2 = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & -1 & 1 \\ 2 & 1 & 2 \end{vmatrix} = (-2-1)\hat{i} - (2-2)\hat{j} + (1+2)\hat{k}$$

$$= -3\hat{i} - 0\hat{j} + 3\hat{k}$$

$$\therefore |\vec{b}_1 \times \vec{b}_2| = \sqrt{(-3)^2 + 0^2 + 3^2} = \sqrt{18} = 3\sqrt{2}$$

$$\text{Again } (\vec{a}_2 - \vec{a}_1) \cdot (\vec{b}_1 \times \vec{b}_2) = (\hat{i} - 3\hat{j} - 2\hat{k}) \cdot (-3\hat{i} + 3\hat{k})$$

$$= (1)(-3) + (-3)(0) + (-2)(3) = -9$$

Putting these values in eqn. (i),

$$\text{S.D. } (d) = \frac{|-9|}{3\sqrt{2}} = \frac{9}{3\sqrt{2}} = \frac{3}{\sqrt{2}} = \frac{3\sqrt{2}}{2}$$

15. Find the shortest distance between the lines

$$\frac{x+1}{7} = \frac{y+1}{-6} = \frac{z+1}{1} \quad \text{and} \quad \frac{x-3}{1} = \frac{y-5}{-2} = \frac{z-7}{1}$$

Sol. Equation of one line is $\frac{z+1}{1} = \frac{y+1}{-6} = \frac{x+1}{7}$

Comparing with $\frac{z-z_1}{z_2-z_1} = \frac{y-y_1}{y_2-y_1} = \frac{x-x_1}{x_2-x_1}$

$$b_1 = \frac{x_1}{-x_1},$$

we


have

$$x_1 = -1, y_1 = -1, z_1 = -1; a_1 = 7, b_1 = -6, c_1 = 1$$

$$\therefore \text{vector form of this line is } \vec{r} = a_1 \hat{i} + b_1 \hat{j} + c_1 \hat{k}$$

$$\text{where } a_1 = (x_1, y_1, z_1) = (-1, -1, -1) = -\hat{i} - \hat{j} - \hat{k}$$

$$\text{and } b_1 = a_1 \hat{i} + b_1 \hat{j} + c_1 \hat{k} = 7\hat{i} - 6\hat{j} + \hat{k}$$



$$\frac{z-3}{1} = \frac{y-5}{-2} = \frac{x-7}{1}$$

Equation of second line is

Comparing with $\frac{z-z_2}{a_2} = \frac{y-y_2}{b_2} = \frac{x-x_2}{c_2}$; we have

$$x_2 = 3, y_2 = 5, z_2 = 7; a_2 = 1, b_2 = -2, c_2 = 1$$

∴ vector form of this second line is $\vec{r} = \vec{a}_2 + \mu \vec{b}_2$

where $\vec{a}_2 = (x_2, y_2, z_2) = (3, 5, 7) = 3\hat{i} + 5\hat{j} + 7\hat{k}$

and $\vec{b}_2 = a_2\hat{i} + b_2\hat{j} + c_2\hat{k} = \hat{i} - 2\hat{j} + \hat{k}$
 we know that S.D. between two skew lines is given by

$$d = \frac{|\vec{a}_2 - \vec{a}_1 \cdot (\vec{b}_1 \times \vec{b}_2)|}{|\vec{b}_1 \times \vec{b}_2|} \dots (i)$$

Now $\vec{a}_2 - \vec{a}_1 = 3\hat{i} + 5\hat{j} + 7\hat{k} - (-\hat{i} - \hat{j} - \hat{k})$
 $= 4\hat{i} + 6\hat{j} + 8\hat{k}$

$$\vec{b}_1 \times \vec{b}_2 = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 7 & -6 & 1 \\ 1 & -2 & 1 \end{vmatrix}$$

$$= (-6 + 2)\hat{i} - (7 - 1)\hat{j} + (-14 + 6)\hat{k}$$

$$= -4\hat{i} - 6\hat{j} - 8\hat{k}$$

$$\therefore |\vec{b}_1 \times \vec{b}_2| = \sqrt{(-4)^2 + (-6)^2 + (-8)^2}$$

$$= \sqrt{16 + 36 + 64} = \sqrt{116}$$

again $(\vec{a}_2 - \vec{a}_1) \cdot (\vec{b}_1 \times \vec{b}_2) = 4(-4) + 6(-6) + 8(-8)$
 $= -16 - 36 - 64 = -116$

Putting these values in (i) 

$$\begin{aligned} \text{S.D. } (d) &= \frac{|-116|}{\sqrt{116}} = \frac{116}{\sqrt{116}} = \sqrt{116} \\ &= \sqrt{4 \times 29} = 2\sqrt{29} \end{aligned}$$

16. Find the shortest distance between the lines whose vector equations are

$$\begin{aligned} \vec{r} &= (\hat{i} + 2\hat{j} + 3\hat{k}) + \lambda(\hat{i} - 3\hat{j} + 2\hat{k}) \text{ and} \\ \vec{r} &= 4\hat{i} + 5\hat{j} + 6\hat{k} + \mu(2\hat{i} + 3\hat{j} + \hat{k}). \end{aligned}$$



Sol. Equation of the first line is

$$\vec{r} = (i + 2j + 3k) + \lambda(i - 3j + 2k) = \vec{a}_1 + \lambda \vec{b}_1$$

Comparing, $\vec{a}_1 = i + 2j + 3k$ and $\vec{b}_1 = i - 3j + 2k$

Equation of second line is

$$\vec{r} = (4i + 5j + 6k) + \mu(2i + 3j + k) = \vec{a}_2 + \mu \vec{b}_2$$

Comparing $\vec{a}_2 = 4i + 5j + 6k$ and $\vec{b}_2 = 2i + 3j + k$

We know that length of S.D. between two (skew) lines is

$$\frac{|\vec{a}_2 - \vec{a}_1 \cdot (\vec{b}_1 \times \vec{b}_2)|}{|\vec{b}_1 \times \vec{b}_2|} \quad \dots(i)$$

$$\begin{aligned} \text{Now } \vec{a}_2 - \vec{a}_1 &= 4i + 5j + 6k - (i + 2j + 3k) \\ &= 4i + 5j + 6k - i - 2j - 3k = 3i + 3j + 3k \end{aligned}$$

$$\text{Again } \vec{b}_1 \times \vec{b}_2 = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & -3 & 2 \\ 2 & 3 & 1 \end{vmatrix}$$

Expanding along first row,

$$\vec{b}_1 \times \vec{b}_2 = i(-3 - 6) - j(1 - 4) + k(3 + 6) = -9i + 3j + 9k$$

$$\begin{aligned} \therefore (\vec{a}_2 - \vec{a}_1) \cdot (\vec{b}_1 \times \vec{b}_2) &= 3(-9) + 3(3) + 3(9) \\ &= -27 + 9 + 27 = 9 \end{aligned}$$

$$\begin{aligned} \text{and } |\vec{b}_1 \times \vec{b}_2| &= \sqrt{(-9)^2 + (3)^2 + (9)^2} = \sqrt{81 + 9 + 81} \\ &= \sqrt{171} = \sqrt{9 \times 19} = 3\sqrt{19} \end{aligned}$$

Putting these values in (i),

$$\text{length of shortest distance} = \frac{9}{3\sqrt{19}} = \frac{3}{\sqrt{19}}$$

17. Find the shortest distance between the lines whose vector equations are

$$\vec{r} = (1 - t) \hat{i} + (t - 2) \hat{j} + (3 - 2t) \hat{k} \text{ and}$$

$$\vec{r} = (s + 1) \hat{i} + (2s - 1) \hat{j} - (2s + 1) \hat{k}.$$

Sol. The first line is $\vec{r} = (1 - t) \hat{i} + (t - 2) \hat{j} + (3 - 2t) \hat{k}$



$$\begin{aligned}
 &= \hat{i} - t\hat{i} + t\hat{j} - 2\hat{j} + 3\hat{k} - 2t\hat{k} \\
 &= (\hat{i} - 2\hat{j} + 3\hat{k}) + t(-\hat{i} + \hat{j} - 2\hat{k}) = \mathbf{a}_1 + t\mathbf{b}_1
 \end{aligned}$$

Comparing $\mathbf{a}_1 = \hat{i} - 2\hat{j} + 3\hat{k}$, $\mathbf{b}_1 = -\hat{i} + \hat{j} - 2\hat{k}$

The second line is $\mathbf{r} = (s+1)\hat{i} + (2s-1)\hat{j} - (2s+1)\hat{k}$

$$\begin{aligned}
 &= s\hat{i} + \hat{i} + 2s\hat{j} - \hat{j} - 2s\hat{k} - \hat{k} \\
 &= (\hat{i} - \hat{j} - \hat{k}) + s(\hat{i} + 2\hat{j} - 2\hat{k}) = \mathbf{a}_2 + s\mathbf{b}_2
 \end{aligned}$$

Comparing $\mathbf{a}_2 = \hat{i} - \hat{j} - \hat{k}$, $\mathbf{b}_2 = \hat{i} + 2\hat{j} - 2\hat{k}$

We know that the S.D. between the two (skew) lines is given by

$$d = \frac{|\mathbf{a}_2 - \mathbf{a}_1 \cdot (\mathbf{b}_1 \times \mathbf{b}_2)|}{|\mathbf{b}_1 \times \mathbf{b}_2|} \quad \dots(i)$$

Now $\mathbf{a}_2 - \mathbf{a}_1 = (\hat{i} - \hat{j} - \hat{k}) - (\hat{i} - 2\hat{j} + 3\hat{k}) = \hat{j} - 4\hat{k}$

$$\mathbf{b}_1 \times \mathbf{b}_2 = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ -1 & 1 & -2 \\ 1 & 2 & -2 \end{vmatrix}$$

$$= (-2+4)\hat{i} - (2+2)\hat{j} + (-2-1)\hat{k} = 2\hat{i} - 4\hat{j} - 3\hat{k}$$

$$\therefore |\mathbf{b}_1 \times \mathbf{b}_2| = \sqrt{2^2 + (-4)^2 + (-3)^2} = \sqrt{29}$$

Again $(\mathbf{a}_2 - \mathbf{a}_1) \cdot (\mathbf{b}_1 \times \mathbf{b}_2) = (\hat{j} - 4\hat{k}) \cdot (2\hat{i} - 4\hat{j} - 3\hat{k})$



$$= (1)(-4) + (-4)(-3) = 8$$

Putting these values in eqn. (i),

$$\text{S.D. } (d) = \frac{181}{\sqrt{29}} = \frac{8}{\sqrt{29}}.$$



Exercise 11.3

Note: Formula for question numbers 1 and 2.

If p is the length of perpendicular from the origin to a plane and \hat{m} is a unit normal vector to the plane, then equation of the plane is $\vec{r} \cdot \hat{m} = p$ (where of course p being length is > 0).



1. In each of the following cases, determine the direction cosines of the normal to the plane and the distance from the origin.

(a) $z = 2$

(b) $x + y + z = 1$

(c) $2x + 3y - z = 5$

(d) $5y + 8 = 0$

Sol. (a) Given: Equation of the plane is $z = 2$

Let us first reduce it to vector form $\vec{r} \cdot \vec{n} = d$
where $d > 0$

or $0x + 0y + 1z = 2$ (Here $d = 2 > 0$)

$\Rightarrow (xi + yj + zk) \cdot (0i + 0j + k) = 2$

($\because a_1a + b_1b + c_1c = (a_1\hat{i} + b_1\hat{j} + c_1\hat{k}) \cdot (a_2\hat{i} + b_2\hat{j} + c_2\hat{k})$)

$\Rightarrow \vec{r} \cdot \vec{n} = 2$ where we know that

$\vec{r} = x\hat{i} + y\hat{j} + z\hat{k} =$ (Position vector of point $P(x, y, z)$)

and here $\vec{n} = 0\hat{i} + 0\hat{j} + \hat{k}$

Now let us reduce $\vec{r} \cdot \vec{n} = d$ to $\vec{r} \cdot \vec{n} = p$

Dividing both sides by $|\vec{n}|$, $\frac{\vec{r} \cdot \vec{n}}{|\vec{n}|} = 2$

i.e., $\vec{r} \cdot \hat{n} = 2 = p$ where $\hat{n} = \frac{\vec{n}}{|\vec{n}|} = \frac{0\hat{i} + 0\hat{j} + \hat{k}}{\sqrt{0+0+1}} = 1$

i.e., $\hat{n} = 0\hat{i} + 0\hat{j} + \hat{k}$ and $p = 2$

\therefore By definition, direction cosines of normal to the plane are coefficients of \hat{i} , \hat{j} , \hat{k} in \hat{n} i.e., $0, 0, 1$ and length

of perpendicular from the origin to the plane is $p = 2$.

(b) Given: Equation of the plane is $x + y + z = 1$

$\Rightarrow 1x + 1y + 1z = 1$ (Here $d = 1 > 0$)

$$\Rightarrow (x \mathbf{i} + y \mathbf{j} + z \mathbf{k}) \cdot (\mathbf{i} + \mathbf{j} + \mathbf{k}) = 1$$

$$\text{i.e., } \vec{r} \cdot \vec{n} = 1 \quad \text{where } \vec{n} = \hat{i} + \hat{j} + \hat{k}$$

Dividing both sides by $|\vec{n}| = \sqrt{1^2 + 1^2 + 1^2} = \sqrt{3}$, we have

$$\vec{r} \cdot \frac{\vec{n}}{|\vec{n}|} = \frac{1}{|\vec{n}|}$$



$$\text{i.e., } \vec{r} \cdot \hat{n} = \frac{1}{\sqrt{3}} = p \text{ where } \hat{n} = \frac{\vec{n}}{|\vec{n}|} = \frac{\hat{i} + \hat{j} + \hat{k}}{\sqrt{3}}$$

$$\text{i.e., } \hat{n} = \frac{1}{\sqrt{3}} \hat{i} + \frac{1}{\sqrt{3}} \hat{j} + \frac{1}{\sqrt{3}} \hat{k} \text{ and } p = \frac{1}{\sqrt{3}}$$

By definition, direction cosines of the normal to the plane are the coefficients of \hat{i} , \hat{j} , \hat{k} in \hat{n} i.e., $\frac{1}{\sqrt{3}}$, $\frac{1}{\sqrt{3}}$,

$\frac{1}{\sqrt{3}}$ and length of perpendicular from the origin to the plane

$$\text{is } p = \frac{1}{\sqrt{3}}.$$

- (c) **Given:** Equation of the plane is $2x + 3y - z = 5$
 $\Rightarrow 2x + 3y + (-1)z = 5$ (Here $d = 5 > 0$)

$$\Rightarrow (x\hat{i} + y\hat{j} + z\hat{k}) \cdot (2\hat{i} + 3\hat{j} - \hat{k}) = 5$$

$$\text{i.e., } \vec{r} \cdot \vec{n} = 5 \text{ where } \vec{n} = 2\hat{i} + 3\hat{j} - \hat{k}$$

Dividing both sides by $|\vec{n}| = \sqrt{4+9+1} = \sqrt{14}$,

$$\text{we have } \vec{r} \cdot \frac{\vec{n}}{|\vec{n}|} = \frac{5}{|\vec{n}|}$$

$$\text{i.e., } \vec{r} \cdot \hat{n} = \frac{5}{\sqrt{14}} = p \text{ where } \hat{n} = \frac{\vec{n}}{|\vec{n}|} = \frac{2\hat{i} + 3\hat{j} - \hat{k}}{\sqrt{4+9+1} = \sqrt{14}}$$

$$\text{i.e., } \hat{n} = \frac{2}{\sqrt{14}} \hat{i} + \frac{3}{\sqrt{14}} \hat{j} - \frac{1}{\sqrt{14}} \hat{k}$$

By definition, direction cosines of the normal to the plane are coefficients of \hat{i} , \hat{j} , \hat{k} in \hat{n} i.e., $\frac{2}{\sqrt{14}}$, $\frac{3}{\sqrt{14}}$,

$-\frac{1}{\sqrt{14}}$ and length of perpendicular from the origin to the

plane is $\frac{5}{\sqrt{14}}$.

- (d) **Given:** Equation of the plane is

$$5x + 8 = 0 \text{ or } 5x = -8$$

Dividing both sides by -1 to make R.H.S. ($= d$) as positive,
 $-5y = 8$ or $0x + (-5)y + 0z = 8$ [Now $d = 8 > 0$
 $\Rightarrow (x\hat{i} + y\hat{j} + z\hat{k}) \cdot (0\hat{i} - 5\hat{j} + 0\hat{k}) = 8$
i.e., $\vec{r} \cdot \vec{n} = 8$ where $\vec{n} = 0\hat{i} - 5\hat{j} + 0\hat{k}$



Dividing both sides by $|\vec{n}| = \sqrt{0^2 + (-5)^2 + 0^2}$

i.e., $|\vec{n}| = \sqrt{25} = 5$

we have $\vec{r} \cdot \frac{\vec{n}}{|\vec{n}|} = \frac{8}{5}$ i.e., $\vec{r} \cdot \hat{n} = \frac{8}{5} = p$

where $\hat{n} = \frac{\vec{n}}{|\vec{n}|} = \frac{0\hat{i} - 5\hat{j} + 0\hat{k}}{5}$

$= 0\hat{i} - \hat{j} + 0\hat{k} = 0\hat{i} - \hat{j} + 0\hat{k}$ and $p = \frac{8}{5}$.

\therefore By definition, direction cosines of the normal to the plane are coefficients of \hat{i} , \hat{j} , \hat{k} in \hat{n} i.e., 0, -1, 0 and

length of perpendicular from the origin to the plane is $\frac{8}{5}$.

2. Find the vector equation of a plane which is at a distance of 7 units from the origin and normal to the vector

$$3\hat{i} + 5\hat{j} - 6\hat{k}.$$

Sol. Here $\vec{n} = 3\hat{i} + 5\hat{j} - 6\hat{k}$

\therefore The unit vector perpendicular to plane is

$$\hat{n} = \frac{\vec{n}}{|\vec{n}|} = \frac{3\hat{i} + 5\hat{j} - 6\hat{k}}{\sqrt{(3)^2 + (5)^2 + (-6)^2}} = \frac{3\hat{i} + 5\hat{j} - 6\hat{k}}{\sqrt{70}}$$

Also $p = 7$

(given)

Hence, the equation of the required plane is $\vec{r} \cdot \hat{n} = p$

i.e., $\vec{r} \cdot \frac{(3\hat{i} + 5\hat{j} - 6\hat{k})}{\sqrt{70}} = 7$

or $\vec{r} \cdot (3\hat{i} + 5\hat{j} - 6\hat{k}) = 7\sqrt{70}$.

3. Find the Cartesian equation of the following planes:

$$\vec{r} \cdot \hat{n} = p$$

$$(a) \vec{r} \cdot (\hat{i} + \hat{j} - \hat{k}) = 2 \quad (b) \vec{r} \cdot (2\hat{i} + 3\hat{j} - 4\hat{k}) = 1$$

$$(c) \vec{r} \cdot [(s - 2t)\hat{i} + (3 - t)\hat{j} + (2s + t)\hat{k}] = 15.$$

Sol. (a) Vector equation of the plane is

$$\vec{r} \cdot (\hat{i} + \hat{j} - \hat{k}) = 2 \quad \dots(i)$$

Putting $\vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$ in (i) (we know that in 3-D,

\vec{r} is the position vector of any point, $P(x, y, z)$),



Cartesian equation of the plane is

$$\begin{aligned} & (x\hat{i} + y\hat{j} + z\hat{k}) \cdot (\hat{i} + \hat{j} - \hat{k}) = 2 \\ \Rightarrow & x(1) + y(1) + z(-1) = 2 \Rightarrow x + y - z = 2. \end{aligned}$$

(b) We know that \vec{r} is the position vector of any arbitrary point P(x, y, z) on the plane.

$$\begin{aligned} \therefore \vec{r} &= x\hat{i} + y\hat{j} + z\hat{k}, \\ \therefore \vec{r} \cdot (2\hat{i} + 3\hat{j} - 4\hat{k}) &= 1 \text{ (given)} \end{aligned}$$

$$\begin{aligned} \Rightarrow & (x\hat{i} + y\hat{j} + z\hat{k}) \cdot (2\hat{i} + 3\hat{j} - 4\hat{k}) = 1 \\ \Rightarrow & 2x + 3y - 4z = 1 \end{aligned}$$

which is the required Cartesian equation of the plane.

(c) Vector equation of the plane is

$$\vec{r} \cdot [(s - 2t)\hat{i} + (3 - t)\hat{j} + (2s + t)\hat{k}] = 15 \quad \dots(i)$$

We know that \vec{r} is the position vector of any point P(x, y, z) on plane (i).

$$\therefore \vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$$

Putting $\vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$ in (i), Cartesian equation of the required plane is

$$(x\hat{i} + y\hat{j} + z\hat{k}) \cdot [(s - 2t)\hat{i} + (3 - t)\hat{j} + (2s + t)\hat{k}] = 15$$

$$\text{i.e., } x(s - 2t) + y(3 - t) + z(2s + t) = 15.$$

4. In the following cases, find the coordinates of the foot of the perpendicular drawn from the origin.

(a) $2x + 3y + 4z - 12 = 0$

(b) $3y + 4z - 6 = 0$

(c) $x + y + z = 1$

$0(0, 0, 0)$

(d) $5y + 8 = 0.$

Sol. (a) Given: Equation of the plane is

$$2x + 3y + 4z - 12 = 0 \quad \dots(i)$$

Given point is O(0, 0, 0)

Let M be the foot of perpendicular drawn from the origin (0, 0, 0) to plane (i).

\therefore By definition, direction ratios of perpendicular OM to plane (i) are coefficients of x, y, z in (i) i.e., 2, 3, 4 = a, b, c.

\therefore Equations of perpendicular OM are

$$\frac{z - z_1}{2} = \frac{y - y_1}{3} = \frac{x - x_1}{4} = \lambda(\text{say}) \quad \left| \quad \frac{z - z_1}{a} = \frac{y - y_1}{b} = \frac{x - x_1}{c} \right.$$

$$\Rightarrow \frac{z}{2} = \frac{y}{3} = \frac{x}{4} = \lambda \quad \Rightarrow \frac{z}{2} = \lambda, \frac{y}{3} = \lambda \text{ and } \frac{x}{4} = \lambda$$

$$\Rightarrow x = 2\lambda, y = 3\lambda, z = 4\lambda$$

\therefore Point M of this line OM is $M(2\lambda, 3\lambda, 4\lambda)$...*(ii)*
for some real λ .



But point M lies on plane (i)

Putting $x = 2\lambda$, $y = 3\lambda$, $z = 4\lambda$ in (i), we have

$$2(2\lambda) + 3(3\lambda) + 4(4\lambda) - 12 = 0$$

$$\Rightarrow 4\lambda + 9\lambda + 16\lambda = 12 \Rightarrow 29\lambda = 12$$

$$\Rightarrow \lambda = \frac{12}{29}$$

Putting $\lambda = \frac{12}{29}$ in (i), foot of perpendicular M $\left(\frac{24}{29}, \frac{36}{29}, \frac{48}{29} \right)$.

(b) For figure, see figure of part (a).

Given: Equation of the plane is $3y + 4z - 6 = 0$... (i)

Given point is $O(0, 0, 0)$

Let M be the foot of perpendicular drawn from the origin to plane (i).

\therefore By definition direction ratios of perpendicular OM to plane (i) are coefficients of x, y, z in (i) i.e., $0, 3, 4 = a, b, c$.

\therefore Equations of perpendicular OM are

$$\frac{z-0}{0} = \frac{y-0}{3} = \frac{x-0}{4} = \lambda(\text{say}) \quad \left| \quad \frac{z-z_1}{a} = \frac{y-y_1}{b} = \frac{x-x_1}{c} \right.$$

$$\Rightarrow \frac{z}{0} = \frac{y}{3} = \frac{x}{4} = \lambda(\text{say}) \Rightarrow \frac{z}{0} = \lambda, \frac{y}{3} = \lambda \text{ and } \frac{x}{4} = \lambda$$

$$\Rightarrow x = 0, y = 3\lambda, z = 4\lambda$$

\therefore Point M of this line OM is $M(0, 3\lambda, 4\lambda)$... (ii)

for some real λ .

But point M lies on plane (i)

Putting $x = 0$, $y = 3\lambda$, $z = 4\lambda$ in (i), we have

$$3(3\lambda) + 4(4\lambda) - 6 = 0 \text{ or } 9\lambda + 16\lambda = 6$$

$$\Rightarrow 25\lambda = 6 \Rightarrow \lambda = \frac{6}{25}$$

Putting $\lambda = \frac{6}{25}$ in (ii), the required foot M of perpendicular

$$\text{is } \left(0, \frac{18}{25}, \frac{24}{25} \right).$$

(c) For figure, see figure of part (a).

Given: Equation of the plane is

$$x + y + z = 1 \quad \dots (i)$$

Given point is $O(0, 0, 0)$

Let M be the foot of perpendicular drawn from the origin $(0, 0, 0)$ to plane (i).

\therefore By definition direction ratios of perpendicular OM to plane (i) are coefficients of x, y, z in (i) i.e., $1, 1, 1 = a, b, c$.

∴ Equations of perpendicular OM are

$$\frac{z-0}{1} = \frac{y-0}{1} = \frac{x-0}{1} \quad \left| \quad \frac{z-z_1}{a} = \frac{y-y_1}{b} = \frac{x-x_1}{c} \right.$$



$$\text{i.e., } \frac{z}{1} = \frac{y}{1} = \frac{x}{1} = \lambda(\text{say}) \quad \therefore \frac{z}{1} = \lambda, \frac{y}{1} = \lambda \text{ and } \frac{x}{1} = \lambda$$

$$\Rightarrow x = \lambda, y = \lambda, z = \lambda$$

\therefore Point M of line OM is $M(\lambda, \lambda, \lambda)$...*(i)*
for some real λ .

But point M lies on plane (i)

Putting $x = \lambda, y = \lambda, z = \lambda$ in (i), we have

$$\lambda + \lambda + \lambda = 1 \Rightarrow 3\lambda = 1 \Rightarrow \lambda = \frac{1}{3}$$

Putting $\lambda = \frac{1}{3}$ in (i), required foot M of perpendicular is

$$\left(\frac{1}{3}, \frac{1}{3}, \frac{1}{3} \right).$$

(d) For figure, see figure of part (a).

Given: Equation of the plane is

$$5y + 8 = 0 \quad \dots(i)$$

Given point is $O(0, 0, 0)$

Let M be the foot of perpendicular drawn from the origin $(0, 0, 0)$ to plane (i).

\therefore By definition, direction ratios of perpendicular OM to plane (i) are coefficients of x, y, z in (i) i.e., $0, 5, 0 = a, b, c$.

\therefore Equations of perpendicular OM are

$$\frac{z-0}{0} = \frac{y-0}{5} = \frac{x-0}{0} \quad \left| \quad \frac{z-z_1}{a} = \frac{y-y_1}{b} = \frac{x-x_1}{c} \right.$$

$$\text{i.e., } \frac{z}{0} = \frac{y}{5} = \frac{x}{0} = \lambda(\text{say}) \quad \therefore \frac{z}{0} = \lambda, \frac{y}{5} = \lambda \text{ and } \frac{x}{0} = \lambda$$

$$\Rightarrow x = 0, y = 5\lambda, z = 0$$

\therefore Point M of line OM is $M(0, 5\lambda, 0)$...*(ii)*
for some real λ .

But point M lies on plane (i)

Putting $x = 0, y = 5\lambda$ and $z = 0$ in (i), we have

$$5(5\lambda) + 8 = 0 \text{ or } 25\lambda = -8$$

$$\Rightarrow \lambda = -\frac{8}{25}$$

Putting $\lambda = -\frac{8}{25}$ in (i), required foot M of perpendicular is

$$\left(0, -\frac{40}{25}, 0 \right) = \left(0, -\frac{8}{5}, 0 \right).$$

$$\left(0, -\frac{8}{5}, 0 \right)$$

5. Find the vector and cartesian equations of the planes
- (a) that passes through the point $(1, 0, -2)$ and the normal vector to the plane is $\hat{i} + \hat{j} - \hat{k}$.
- (b) that passes through the point $(1, 4, 6)$ and the normal vector to the plane is $\hat{i} - 2\hat{j} + \hat{k}$.



Sol. (a) *Vector form of equation of the plane*

The given point on the plane is $(1, 0, -2)$

∴ The position vector of the given point is

$$\vec{a} = (1, 0, -2) = \hat{i} + 0\hat{j} - 2\hat{k} = \hat{i} - 2\hat{k}$$

Also Given: Normal vector to the plane is

$$\vec{n} = \hat{i} + \hat{j} - \hat{k}$$

∴ Vector equation of the required plane is

$$(\vec{r} - \vec{a}) \cdot \vec{n} = 0 \quad \text{i.e.,} \quad \vec{r} \cdot \vec{n} - \vec{a} \cdot \vec{n} = 0$$

$$\text{i.e.,} \quad \vec{r} \cdot \vec{n} = \vec{a} \cdot \vec{n}$$

Putting values of \vec{a} and \vec{n} ,

$$\vec{r} \cdot (\hat{i} + \hat{j} - \hat{k}) = (\hat{i} - 2\hat{k}) \cdot (\hat{i} + \hat{j} - \hat{k})$$

$$\text{i.e.,} \quad \vec{r} \cdot (\hat{i} + \hat{j} - \hat{k}) = 1(1) + 0(1) + (-2)(-1) = 1 + 2 = 3$$

$$\text{i.e.,} \quad \vec{r} \cdot (\hat{i} + \hat{j} - \hat{k}) = 3$$

Cartesian form of equation of the plane

The plane passes through the point $(1, 0, -2) = (x_1, y_1, z_1)$

Normal vector to the plane is $\vec{n} = \hat{i} + \hat{j} - \hat{k}$

∴ Direction ratios of normal to the plane are coefficients

of \hat{i} , \hat{j} , \hat{k} in \vec{n} i.e., $1, 1, -1$.

∴ Cartesian equation of the required plane is

$$a(x - x_1) + b(y - y_1) + c(z - z_1) = 0$$

$$\text{or} \quad 1(x - 1) + 1(y - 0) - (z + 2) = 0$$

$$\text{i.e.,} \quad x - 1 + y - z - 2 = 0$$

$$\text{i.e.,} \quad x + y - z = 3.$$

(b) **Vector form of the equation of the plane**

The given point on the plane is $(1, 4, 6)$.

∴ The position vector of the given point is

$$\vec{a} = (1, 4, 6) = \hat{i} + 4\hat{j} + 6\hat{k}$$

Also Given: normal vector to the plane is $\vec{n} = \hat{i} - 2\hat{j} + \hat{k}$.

∴ Equation of the plane is $(\vec{r} - \vec{a}) \cdot \vec{n} = 0$

$$\text{or} \quad \vec{r} \cdot \vec{n} - \vec{a} \cdot \vec{n} = 0 \quad \text{i.e.,} \quad \vec{r} \cdot \vec{n} = \vec{a} \cdot \vec{n}$$

Putting values of \vec{a} and \vec{n}

$$\begin{aligned} \vec{r} \cdot (\hat{i} - 2\hat{j} + \hat{k}) &= (\hat{i} + 4\hat{j} + 6\hat{k}) \cdot (\hat{i} - 2\hat{j} + \hat{k}) \\ &= 1 - 8 + 6 = -1 \end{aligned} \quad \dots(i)$$

Cartesian Form

The plane passes through the point $(1, 4, 6) = (x_1, y_1, z_1)$.

Normal vector to the plane is $\vec{n} = \hat{i} - 2\hat{j} + \hat{k}$.



\therefore D.R.'s of the normal to the plane are coefficients of \hat{i} , \hat{j} , \hat{k} in \vec{n}

i.e., $1, -2, 1 = a, b, c$

\therefore Equation of the required plane is

$$\text{or } a(x - x_1) + b(y - y_1) + c(z - z_1) = 0$$

$$\text{or } 1(x - 1) - 2(y - 4) + 1(z - 6) = 0$$

$$\text{or } x - 1 - 2y + 8 + z - 6 = 0$$

$$\text{or } x - 2y + z + 1 = 0$$

Alternatively for Cartesian form

$$\text{From eqn. (i), } (x\hat{i} + y\hat{j} + z\hat{k}) \cdot (\hat{i} - 2\hat{j} + \hat{k}) = -1$$

$$\text{or } x - 2y + z = -1 \text{ or } x - 2y + z + 1 = 0.$$

6. Find the equations of the planes that passes through three points:

(a) $(1, 1, -1), (6, 4, -5), (-4, -2, 3)$

(b) $(1, 1, 0), (1, 2, 1), (-2, 2, -1)$

Sol. We know that through three collinear points A, B, C i.e., through a straight line, we can pass an infinite number of planes.

(a) The three given points are

$$A(1, 1, -1), B(6, 4, -5), C(-4, -2, 3)$$

Let us examine whether these points are collinear.

Direction ratios of line AB are

$$6 - 1, 4 - 1, -5 + 1 \quad | \quad x_2 - x_1, y_2 - y_1, z_2 - z_1$$

$$= 5, 3, -4 = a_1, b_1, c_1$$

Again direction ratios of line BC are

$$-4 - 6, -2 - 4, 3 - (-5) = -10, -6, 8 = a_2, b_2, c_2$$

$$\text{Here } \frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2} \Rightarrow \frac{5}{-10} = \frac{3}{-6} = -\frac{4}{8}$$

$$\Rightarrow -\frac{1}{2} = -\frac{1}{2} = -\frac{1}{2} \text{ which is true.}$$

\therefore Lines AB and BC are parallel.

But B is their common point.

\therefore Points A, B and C are collinear and hence an infinite number of planes can be drawn through the three given collinear points.

(b) The three given points are

$$A(1, 1, 0) = (x_1, y_1, z_1), B(1, 2, 1) = (x_2, y_2, z_2)$$

$$\text{and } C(-2, 2, -1) = (x_3, y_3, z_3)$$

Let us examine whether these points are collinear.

Direction ratios of line AB are

$$1 - 1, 2 - 1, 1 - 0 \quad | \quad x_2 - x_1, y_2 - y_1, z_2 - z_1$$

$$\text{i.e., } 0, 1, 1 = a_1, b_1, c_1$$

Direction ratios of line BC are

$$-2 - 1, 2 - 2, -1 - 1 = -3, 0, -2 = a_2, b_2, c_2$$

$$\text{Here } \frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2} =$$

$$= \frac{c_1 c_2}{\vec{1}} \Rightarrow \frac{0}{\frac{1}{-3}} =$$
$$0$$
$$-2$$

which is not true.



\therefore Points A, B, C are not collinear.

\therefore Equation of the unique plane passing through these three points A, B, C is

$$\begin{vmatrix} x - x_1 & y - y_1 & z - z_1 \\ x_2 - x_1 & y_2 - y_1 & z_2 - z_1 \\ x_3 - x_1 & y_3 - y_1 & z_3 - z_1 \end{vmatrix} = 0$$

$$\Rightarrow \begin{vmatrix} z - 1 & y - 1 & x - 0 \\ 1 - 1 & 2 - 1 & 1 - 0 \\ -2 - 1 & 2 - 1 & -1 - 0 \end{vmatrix} = 0 \Rightarrow \begin{vmatrix} z - 1 & y - 1 & x \\ 0 & 1 & 1 \\ -3 & 1 & -1 \end{vmatrix} = 0$$

Expanding along first row,

$$\begin{aligned} & (x - 1)(-1 - 1) - (y - 1)(0 + 3) + z(0 + 3) = 0 \\ \Rightarrow & -2(x - 1) - 3(y - 1) + 3z = 0 \\ \Rightarrow & -2x + 2 - 3y + 3 + 3z = 0 \\ \Rightarrow & -2x - 3y + 3z + 5 = 0 \\ \Rightarrow & 2x + 3y - 3z - 5 = 0 \\ \text{or} & 2x + 3y - 3z = 5 \end{aligned}$$

which is the equation of required plane.

7. Find the intercepts cut off by the plane $2x + y - z = 5$.

Sol. Equation of the plane is $2x + y - z = 5$

Dividing every term by 5, (to make R.H.S. 1)

$$\frac{2z}{5} + \frac{y}{5} - \frac{x}{5} = 1 \text{ or } \frac{z}{\left(\frac{5}{2}\right)} + \frac{y}{5} + \frac{x}{-5} = 1$$

Comparing with intercept form $\frac{z}{a} + \frac{y}{b} + \frac{x}{c} = 1$, we have

$a = \frac{5}{2}$, $b = 5$, $c = -5$ which are the intercepts cut off by the plane on x -axis, y -axis and z -axis respectively.

8. Find the equation of the plane with intercept 3 on the y -axis and parallel to ZOX plane.

Sol. We know that equation of ZOX plane is $y = 0$.

\therefore Equation of any plane parallel to ZOX plane is $y = k$... (i)

(\because Equation of any plane parallel to the plane

$ax + by + cz + d = 0$ is $ax + by + cz + k = 0$

i.e., change only the constant term)

To find k . Plane (i) makes an intercept 3 on the y -axis ($\Rightarrow x = 0$ and $z = 0$) *i.e.*, plane (i) passes through $(0, 3, 0)$.

Putting $x = 0$, $y = 3$ and $z = 0$ in (i), $3 = k$.

Putting $k = 3$ in (i), equation of required plane is $y = 3$.

9. Find the equation of the plane through the intersection of the planes $3x - y + 2z - 4 = 0$ and $x + y + z - 2 = 0$ and the point $(2, 2, 1)$.

Sol. Equations of the given planes are

$$3x - y + 2z - 4 = 0 \quad \text{and} \quad x + y + z - 2 = 0$$



(Here R.H.S. of each equation is already zero)

We know that equation of any plane through the intersection of these two planes is

$$\text{L.H.S. of plane I} + \lambda(\text{L.H.S. of plane II}) = 0$$

$$\text{i.e., } 3x - y + 2z - 4 + \lambda(x + y + z - 2) = 0 \quad \dots(i)$$

To find λ . Given: Required plane (i) passes through the point(2, 2, 1).

Putting $x = 2, y = 2$ and $z = 1$ in (i),

$$6 - 2 + 2 - 4 + \lambda(2 + 2 + 1 - 2) = 0$$

$$\text{or } 2 + 3\lambda = 0 \Rightarrow 3\lambda = -2 \Rightarrow \lambda = -\frac{2}{3}$$

Putting $\lambda = -\frac{2}{3}$ in (i), equation of required plane is

$$3x - y + 2z - 4 - \frac{2}{3}(x + y + z - 2) = 0$$

$$\Rightarrow 9x - 3y + 6z - 12 - 2x - 2y - 2z + 4 = 0$$

$$\Rightarrow 7x - 5y + 4z - 8 = 0.$$

10. Find the vector equation of the plane passing through the

intersection of the planes $\vec{r} \cdot (2\hat{i} + 2\hat{j} - 3\hat{k}) = 7,$

$\vec{r} \cdot (2\hat{i} + 5\hat{j} + 3\hat{k}) = 9$ and through the point (2, 1, 3).

Sol. Vector equation of first plane is

$$\vec{r} \cdot (2\hat{i} + 2\hat{j} - 3\hat{k}) = 7 \text{ i.e. } (x\hat{i} + y\hat{j} + z\hat{k}) \cdot (2\hat{i} + 2\hat{j} - 3\hat{k}) = 7$$

$$\text{i.e. } 2x + 2y - 3z - 7 = 0 \quad (\text{making R.H.S. zero}) \quad \dots(i)$$

Vector equation of second plane is

$$\vec{r} \cdot (2\hat{i} + 5\hat{j} + 3\hat{k}) = 9 \text{ i.e. } (x\hat{i} + y\hat{j} + z\hat{k}) \cdot (2\hat{i} + 5\hat{j} + 3\hat{k}) = 9$$

$$\text{i.e. } 2x + 5y + 3z - 9 = 0 \quad (\text{making R.H.S. zero}) \quad \dots(ii)$$

We know that equation of any plane passing through the line of intersection of planes (i) and (ii) is

$$\text{L.H.S of (i)} + \lambda \text{ L.H.S of (ii)} = 0$$

$$\text{i.e. } 2x + 2y - 3z - 7 + \lambda (2x + 5y + 3z - 9) = 0$$

$$\text{i.e. } 2x + 2y - 3z - 7 + 2\lambda x + 5\lambda y + 3\lambda z - 9\lambda = 0$$

$$\text{i.e. } (2 + 2\lambda)x + (2 + 5\lambda)y + (-3 + 3\lambda)z = 7 + 9\lambda \quad \dots(iii)$$

To find λ : Given plane passes through the point (2,1,3) putting $x = 2, y = 1, z = 3$

$$(2 + 2\lambda) 2 + (2 + 5\lambda) 1 + (-3 + 3\lambda) 3 = 7 + 9\lambda$$

$$\text{or } 4 + 4\lambda + 2 + 5\lambda - 9 + 9\lambda = 7 + 9\lambda$$

$$9\lambda - 3 = 7 \Rightarrow 9\lambda = 10 \Rightarrow \lambda = \frac{10}{9}$$



Putting $\lambda = \frac{10}{9}$ in (iii), equation of required plane is

$$\left(2 + \frac{20}{9}\right)x + \left(2 + \frac{50}{9}\right)y + \left(-3 + \frac{30}{9}\right)z = 7 + 10$$

$$\text{or } \frac{38}{9}x + \frac{68}{9}y + \frac{3}{9}z = 17$$

Multiplying by L.C.M. = 9, $38x + 68y + 3z = 153$

$$\text{or } x(38) + y(68) + z(3) = 153$$

$$\text{or } (x\hat{i} + y\hat{j} + z\hat{k}) \cdot (38\hat{i} + 68\hat{j} + 3\hat{k}) = 153$$

$$\text{i.e. } \vec{r} \cdot (38\hat{i} + 68\hat{j} + 3\hat{k}) = 153$$

which is the required vector equation of the plane.

11. Find the equation of the plane through the line of intersection of the planes $x + y + z = 1$ and $2x + 3y + 4z = 5$ which is perpendicular to the plane $x - y + z = 0$.

Sol. Equations of the given planes are

$$x + y + z = 1 \quad \text{and} \quad 2x + 3y + 4z = 5$$

Making R.H.S. zero, equations of the planes are

$$x + y + z - 1 = 0 \quad \text{and} \quad 2x + 3y + 4z - 5 = 0.$$

We know that equation of any plane through the intersection of the two planes is

$$(\text{L.H.S. of I}) + \lambda(\text{L.H.S. of II}) = 0$$

$$\text{i.e., } x + y + z - 1 + \lambda(2x + 3y + 4z - 5) = 0 \quad \dots(i)$$

$$\text{i.e., } x + y + z - 1 + 2\lambda x + 3\lambda y + 4\lambda z - 5\lambda = 0$$

$$\text{i.e., } (1 + 2\lambda)x + (1 + 3\lambda)y + (1 + 4\lambda)z - 1 - 5\lambda = 0$$

Given: This plane is perpendicular to the plane

$$x - y + z = 0$$

$$\therefore \mathbf{a_1 a_2 + b_1 b_2 + c_1 c_2 = 0}$$

i.e., Product of coefficients of $x + \dots = 0$

$$\therefore (1 + 2\lambda) - (1 + 3\lambda) + 1 + 4\lambda = 0$$

$$\Rightarrow 1 + 2\lambda - 1 - 3\lambda + 1 + 4\lambda = 0 \Rightarrow 3\lambda + 1 = 0 \Rightarrow 3\lambda = -1$$

$$\Rightarrow \lambda = \frac{-1}{3}$$

Putting $\lambda = \frac{-1}{3}$ in (i), equation of required plane is

$$x + y + z - 1 - \frac{1}{3}(2x + 3y + 4z - 5) = 0$$

Multiplying by L.C.M.  **CUET Academy**

$$3x + 3y + 3z - 3 - 2x - 3y - 4z + 5 = 0 \Rightarrow x - z + 2 = 0.$$

12. Find the angle between the planes whose vector equations are

$$\vec{r} \cdot (2\hat{i} + 2\hat{j} - 3\hat{k}) = 5 \text{ and } \vec{r} \cdot (3\hat{i} - 3\hat{j} + 5\hat{k}) = 3.$$



Sol. Equation of one plane is

$$\vec{r} \cdot (2\hat{i} + 2\hat{j} - 3\hat{k}) = 5 \quad \dots(i)$$

Comparing (i) with $\vec{r} \cdot \vec{n}_1 = d_1$, we have

$$\text{normal vector to plane (i) is } \vec{n}_1 = 2\hat{i} + 2\hat{j} - 3\hat{k}$$

$$\text{Equation of second plane is } \vec{r} \cdot (3\hat{i} - 3\hat{j} + 5\hat{k}) = 3 \quad \dots(ii)$$

Comparing (ii) with $\vec{r} \cdot \vec{n}_2 = d_2$, we have

$$\text{normal vector to plane (ii) is } \vec{n}_2 = 3\hat{i} - 3\hat{j} + 5\hat{k}$$

Let θ be the acute angle between planes (i) and (ii).

\therefore By definition, angle between normals \vec{n}_1 and \vec{n}_2 to planes (i) and (ii) is also θ .

$$\begin{aligned} \therefore \cos \theta &= \frac{\vec{n}_1 \cdot \vec{n}_2}{|\vec{n}_1| |\vec{n}_2|} = \frac{12(3) + 2(-3) + (-3)5I}{\sqrt{4+4+9} \sqrt{9+9+25}} \\ &= \frac{16 - 6 - 15I}{\sqrt{17} \sqrt{43}} = \frac{1 - 15I}{\sqrt{17} \times 43} = \frac{15}{\sqrt{731}} \quad \therefore \theta = \cos^{-1} \frac{15}{\sqrt{731}} \end{aligned}$$

13. In the following cases, determine whether the given planes are parallel or perpendicular and in case they are neither, find the angle between them.

(a) $7x + 5y + 6z + 30 = 0$ and $3x - y - 10z + 4 = 0$

(b) $2x + y + 3z - 2 = 0$ and $x - 2y + 5 = 0$

(c) $2x - 2y + 4z + 5 = 0$ and $3x - 3y + 6z - 1 = 0$

(d) $2x - y + 3z - 1 = 0$ and $2x - y + 3z + 3 = 0$

(e) $4x + 8y + z - 8 = 0$ and $y + z - 4 = 0$.

Sol. (a) Equations of the given planes are

$$7x + 5y + 6z + 30 = 0$$

$$(a_1x + b_1y + c_1z + d_1 = 0)$$

and $3x - y - 10z + 4 = 0$ ($a_2x + b_2y + c_2z + d_2 = 0$)

Here $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$ becomes $\frac{7}{3} = \frac{5}{-1} = \frac{6}{-10}$ which is

not true.

\therefore The two planes are not parallel.

Again $a_1a_2 + b_1b_2 + c_1c_2 = 21 - 5 - 60 = 21 - 65 = -44 \neq 0$

\therefore Planes are not perpendicular.

Now let θ be the angle between the two planes.

$$\begin{aligned} \therefore \cos \theta &= \frac{|a_1a_2 + b_1b_2 + c_1c_2|}{\sqrt{a_1^2 + b_1^2 + c_1^2} \sqrt{a_2^2 + b_2^2 + c_2^2}} \\ &= \frac{|7(3) + 5(-1) + 6(-10)|}{\sqrt{(7)^2 + (5)^2 + (6)^2} \sqrt{(3)^2 + (-1)^2 + (-10)^2}} \end{aligned}$$



$$= \frac{121 - 5 - 60I}{\sqrt{49 + 25 + 36} \sqrt{9 + 1 + 100}} = \frac{I - 44I}{\sqrt{110} \sqrt{110}}$$

$$= \frac{I - 44I}{110} = \frac{44}{110} = \frac{2}{5} \quad \therefore \theta = \cos^{-1} \left(\frac{2}{5} \right)$$

- (b) Equations of the given planes are

$$2x + y + 3z - 2 = 0 \quad (a_1x + b_1y + c_1z + d_1 = 0)$$

$$\text{and } x - 2y + 5 = 0 \quad \text{i.e., } x - 2y + 0z + 5 = 0$$

$$(a_2x + b_2y + c_2z + d_2 = 0)$$

Are these planes parallel?

Here $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2} \Rightarrow \frac{2}{1} = \frac{1}{-2} = \frac{3}{0}$ which is not true.

$$\begin{array}{ccc} a_2 & b_2 & c_2 \\ \downarrow & & \end{array} \quad \begin{array}{ccc} 1 & -2 & 0 \end{array}$$

(Ratio of coefficients of x in equations of two planes) \therefore The given planes are not parallel.**Are these planes perpendicular?**

$$a_1a_2 + b_1b_2 + c_1c_2 = 2(1) + 1(-2) + 3(0) = 2 - 2 + 0 = 0$$

(Product of coefficients of x) \therefore Given planes are perpendicular.

- (c) Equations of the given planes are

$$2x - 2y + 4z + 5 = 0 \quad (a_1x + b_1y + c_1z + d_1 = 0) \text{ and}$$

$$3x - 3y + 6z - 1 = 0 \quad (a_2x + b_2y + c_2z + d_2 = 0)$$

Are these planes parallel?

Here $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2} \Rightarrow \frac{2}{3} = \frac{-2}{-3} = \frac{4}{6} \Rightarrow \frac{2}{3} = \frac{2}{3} = \frac{2}{3}$

which is true.

 \therefore The given planes are parallel.

- (d) Equations of the given planes are

$$2x - y + 3z - 1 = 0 \quad (a_1x + b_1y + c_1z + d_1 = 0) \text{ and}$$

$$2x - y + 3z + 3 = 0 \quad (a_2x + b_2y + c_2z + d_2 = 0)$$

Are these planes parallel?

Here $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2} \Rightarrow \frac{2}{2} = \frac{-1}{-1} = \frac{3}{3} \Rightarrow 1 = 1 = 1$

which is true.

 \therefore The given planes are parallel.

- (e) Equations of the given planes are

$$4x + 8y + z + 1 = 0 \quad (a_1x + b_1y + c_1z + d_1 = 0)$$

and $y + z - 4 = 0$ i.e., $0x + y + z - 4 = 0$
($a_2x + b_2y + c_2z + d_2 = 0$)

Are these planes parallel?

Here $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2} \Rightarrow \frac{4}{0} = \frac{8}{1} = \frac{1}{1}$ which is not true.



∴ The given planes are not parallel.

Are these planes perpendicular?

$$\text{Here } a_1a_2 + b_1b_2 + c_1c_2 = 4(0) + 8(1) + 1(1) \\ = 0 + 8 + 1 = 9 \neq 0 \quad \dots(i)$$

∴ The given planes are not perpendicular.

To find the (acute) angle θ between the given planes.

$$\begin{aligned} \therefore \cos \theta &= \frac{a_1a_2 + b_1b_2 + c_1c_2}{\sqrt{a_1^2 + b_1^2 + c_1^2} \sqrt{a_2^2 + b_2^2 + c_2^2}} = \frac{\vec{n}_1 \cdot \vec{n}_2}{|\vec{n}_1| |\vec{n}_2|} \\ &= \frac{4(0) + 8(1) + 1(1)}{\sqrt{16 + 64 + 1} \sqrt{0^2 + 1^2 + 1^2}} = \frac{18 + 1}{\sqrt{81} \sqrt{2}} \\ &= \frac{9}{9\sqrt{2}} = \frac{1}{\sqrt{2}} = \cos 45^\circ \quad \therefore \theta = 45^\circ. \end{aligned}$$

14. In the following cases find the distances of each of the given points from the corresponding given plane.

Point	Plane
(a) (0, 0, 0)	$3x - 4y + 12z = 3$
(b) (3, -2, 1)	$2x - y + 2z + 3 = 0$
(c) (2, 3, -5)	$x + 2y - 2z = 9$
(d) (-6, 0, 0)	$2x - 3y + 6z - 2 = 0.$

Sol. (a) Distance (of course perpendicular) of the point (0, 0, 0) from the plane $3x - 4y + 12z = 3$ or $3x - 4y + 12z - 3 = 0$ (Making R.H.S. zero) is

$$\begin{aligned} \frac{|ax_1 + by_1 + cz_1 + d|}{\sqrt{a^2 + b^2 + c^2}} &= \frac{|3(0) - 4(0) + 12(0) - 3|}{\sqrt{(3)^2 + (-4)^2 + (12)^2}} \\ &= \frac{|-3|}{\sqrt{9 + 16 + 144}} = \frac{3}{\sqrt{169}} = \frac{3}{13}. \end{aligned}$$

- (b) Length of perpendicular from the point (3, -2, 1) on the plane $2x - y + 2z + 3 = 0$ (Substitute the point for x, y, z in L.H.S. of Eqn. of plane and divide by $\sqrt{a^2 + b^2 + c^2}$)

$$= \frac{|2(3) - (-2) + 2(1) + 3|}{\sqrt{(2)^2 + (-1)^2 + (2)^2}} = \frac{|16 + 2 + 2 + 3|}{\sqrt{4 + 1 + 4}} = \frac{13}{\sqrt{9}} = \frac{13}{3}$$

- (c) Length of perpendicular from the point (2, 3, -5) on the plane $x + 2y - 2z = 9$ or $x + 2y - 2z - 9 = 0$ (Making R.H.S. zero)

$$= \frac{|2 + 2(3) - 2(-5) - 9|}{\sqrt{(1)^2 + (2)^2 + (-2)^2}} = \frac{|12 + 6 + 10 - 9|}{\sqrt{1 + 4 + 4}} = \frac{9}{\sqrt{9}} = \frac{9}{3} = 3.$$

- (d) Distance of the point (-6, 0, 0) from the plane

$$2x - 3y + 6z - 2 = 0$$

(Here R.H.S. is already zero)



$$\begin{aligned} &= \frac{Iax_1 + by_1 + cz_1 + dI}{\sqrt{a^2 + b^2 + c^2}} = \frac{I2(-6) - 3(0) + 6(0) - 2I}{\sqrt{(2)^2 + (-3)^2 + (6)^2}} \\ &= \frac{I-12 - 2I}{\sqrt{4+9+36}} = \frac{I-14I}{\sqrt{49}} = \frac{14}{7} = 2. \end{aligned}$$



MISCELLANEOUS EXERCISE

1. Show that the line joining the origin to the point $(2, 1, 1)$ is perpendicular to the line determined by the points $(3, 5, -1)$, $(4, 3, -1)$.

Sol. We know that direction ratios of the line joining the origin $(0, 0, 0)$ to the point $(2, 1, 1)$ are

$$\begin{aligned}x_2 - x_1, y_2 - y_1, z_2 - z_1 &= 2 - 0, 1 - 0, 1 - 0 \\ &= 2, 1, 1 = a_1, b_1, c_1.\end{aligned}$$

Similarly, direction ratios of the line joining the points $(3, 5, -1)$ and $(4, 3, -1)$ are

$$4 - 3, 3 - 5, -1 - (-1) = 1, -2, 0 = a_2, b_2, c_2.$$


For these two lines $a_1a_2 + b_1b_2 + c_1c_2$

$$= 2(1) + 1(-2) + 1(0) = 2 - 2 + 0 = 0$$

Therefore, the two given lines are perpendicular to each other.

2. If l_1, m_1, n_1 and l_2, m_2, n_2 are the direction cosines of two mutually perpendicular lines, show that the direction cosines of the line perpendicular to both of these are $m_1n_2 - m_2n_1, n_1l_2 - n_2l_1, l_1m_2 - l_2m_1$.

Sol.



l_1, m_1, n_1 ; and l_2, m_2, n_2 are d.c.'s of two mutually perpendicular given lines L_1 and L_2 (say).

Let \hat{n}_1 and \hat{n}_2 be the unit vectors along these lines L_1 and L_2 .

$$\therefore \hat{n}_1 = l_1 \hat{i} + m_1 \hat{j} + n_1 \hat{k} \quad \text{and} \quad \hat{n}_2 = l_2 \hat{i} + m_2 \hat{j} + n_2 \hat{k}$$

Let L be the line \perp to both the lines L_1 and L_2 . Let \hat{n} be a unit vector along line $L \perp$ to both lines L_1 and L_2

$$\therefore \hat{n} = \hat{n}_1 \times \hat{n}_2$$



$$n = \frac{\hat{n}_1 \times \hat{n}_2}{|\hat{n}_1 \times \hat{n}_2|} = \frac{\hat{n}_1 \cdot \hat{n}_2}{|\hat{n}_1| |\hat{n}_2| \sin 90^\circ}$$



$$\Rightarrow \vec{n} = \vec{n}_1 \times \vec{n}_2 = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ l_1 & m_1 & n_1 \\ l_2 & m_2 & n_2 \end{vmatrix}$$

$$\text{or } \vec{n} = (m_1 n_2 - m_2 n_1) \hat{i} - (l_1 n_2 - l_2 n_1) \hat{j} + (l_1 m_2 - l_2 m_1) \hat{k}$$

Now because \vec{n} is a unit vector, therefore its components are its direction cosines.

Thus d.c.'s of \vec{n} are $\frac{m_1 n_2 - m_2 n_1}{|\vec{n}|}$, $\frac{l_2 n_1 - l_1 n_2}{|\vec{n}|}$, $\frac{l_1 m_2 - l_2 m_1}{|\vec{n}|}$
 i.e., d.c.'s of line L are $\frac{m_1 n_2 - m_2 n_1}{|\vec{n}|}$, $\frac{l_2 n_1 - l_1 n_2}{|\vec{n}|}$, $\frac{l_1 m_2 - l_2 m_1}{|\vec{n}|}$.

3. Find the angle between the lines whose direction ratios are a, b, c and b - c, c - a, a - b.

Sol. Given: Direction ratios of one line are a, b, c

$$\Rightarrow \text{A vector along this line is } \vec{b}_1 = a \hat{i} + b \hat{j} + c \hat{k}$$

Given: Direction ratios of second line are b - c, c - a, a - b

\Rightarrow A vector along the second line is

$$\vec{b}_2 = (b - c) \hat{i} + (c - a) \hat{j} + (a - b) \hat{k}$$

Let θ be the angle between the two lines.

$$\begin{aligned} \text{We know that } \cos \theta &= \frac{\vec{b}_1 \cdot \vec{b}_2}{|\vec{b}_1| |\vec{b}_2|} \\ &= \frac{a(b - c) + b(c - a) + c(a - b)}{\sqrt{a^2 + b^2 + c^2} \sqrt{(b - c)^2 + (c - a)^2 + (a - b)^2}} \end{aligned}$$

$$\text{or } \cos \theta = \frac{ab - ac + bc - ab + ac - bc}{\sqrt{a^2 + b^2 + c^2} \sqrt{(b - c)^2 + (c - a)^2 + (a - b)^2}}$$


$$\begin{aligned} &= \frac{0}{\sqrt{a^2 + b^2 + c^2} \sqrt{(b - c)^2 + (c - a)^2 + (a - b)^2}} \\ &= 0 = \cos 90^\circ \quad \therefore \theta = 90^\circ. \end{aligned}$$

4. Find the equation of a line parallel to x-axis and passing through the origin.

Sol. We know that a unit vector along x-

$$\text{axis is } \hat{i} = \hat{i} + 0 \hat{j} + 0 \hat{k}$$

\therefore By definition, direction cosines of



$\hat{i}, \hat{j}, \hat{k}$
x-axis are coefficients of $\hat{i}, \hat{j}, \hat{k}$ in
the unit vector *i.e.*, $1, 0, 0 = l, m, n$.
 \therefore Equation of the required line



passing through the origin $(0, 0, 0)$ and parallel to x -axis.
(In fact this required line is x -axis itself)

$$\text{is } \frac{z-0}{1} = \frac{y-0}{0} = \frac{x-0}{0} \quad \text{i.e.,} \quad \frac{z}{1} = \frac{y}{0} = \frac{x}{0}$$

Remark. Whenever it is not mentioned “find vector equation of the line (or plane)”, we should find cartesian equation only.

However vector equation of the required line in the above question

$$\text{is } \vec{r} = \vec{a} + \lambda \vec{b}$$


$$\text{(Here } \vec{a} = \vec{0} \text{ and } \vec{b} = \hat{i} \text{)}$$

$$\text{i.e., } \vec{r} = \vec{0} + \lambda \hat{i} \Rightarrow \vec{r} = \lambda \hat{i}.$$

5. If the coordinates of the points A, B, C, D be $(1, 2, 3)$, $(4, 5, 7)$, $(-4, 3, -6)$ and $(2, 9, 2)$ respectively, then find the angle between the lines AB and CD.

Sol. Given: Points A(1, 2, 3), B(4, 5, 7), C(-4, 3, -6) and D(2, 9, 2).

\therefore Direction ratios of line AB are $x_2 - x_1, y_2 - y_1, z_2 - z_1$



$$\text{i.e., } 4 - 1, 5 - 2, 7 - 3 = 3, 3, 4 = a_1, b_1, c_1$$

$$\therefore \text{ A vector along the line AB is } \vec{b}_1 = 3\hat{i} + 3\hat{j} + 4\hat{k}$$

Similarly direction ratios of line CD are

$$2 - (-4), 9 - 3, 2 - (-6) = 6, 6, 8 = a_2, b_2, c_2.$$

$$\therefore \text{ A vector along the line CD is } \vec{b}_2 = 6\hat{i} + 6\hat{j} + 8\hat{k}$$

Let θ be the angle between the lines AB and CD.

$$\text{We know that } \cos \theta = \frac{Ia_1a_2 + b_1b_2 + c_1c_2I}{\sqrt{a_1^2 + b_1^2 + c_1^2} \sqrt{a_2^2 + b_2^2 + c_2^2}}$$

$$= \frac{Ib_1 \cdot b_2I}{\sqrt{9+9+16} \sqrt{36+36+64}} = \frac{I3(6) + 3(6) + 4(8)I}{\sqrt{44} \sqrt{136}}$$

$$\frac{1b_1 + 1b_2 + 1}{\sqrt{34} \sqrt{136}} = \frac{68}{\sqrt{34 \times 136}} = \frac{68}{\sqrt{34 \times 34 \times 4}} = \frac{68}{34 \times 2}$$



$$= 1 = \cos 0^\circ \quad \therefore \theta = 0$$

\therefore Lines AB and CD are parallel.

6. If the lines $\frac{x-1}{-3} = \frac{y-2}{2k} = \frac{z-3}{2}$ and $\frac{x-1}{3k} = \frac{y-1}{1} =$

$\frac{z-6}{-5}$ are perpendicular, find the value of k .

Sol. Given: Equation of one line is $\frac{z-1}{-3} = \frac{y-2}{2k} = \frac{x-3}{2}$

(It is standard form because coefficient of x, y, z each is unity)

Direction ratios of this line are its denominators

$$-3, 2k, 2 = a_1, b_1, c_1$$

(\Rightarrow a vector along this line is $\vec{b}_1 = -3\hat{i} + 2k\hat{j} + 2\hat{k}$)

Equation of second line is $\frac{z-1}{3k} = \frac{y-1}{1} = \frac{x-6}{-5}$ (Standard form)

Direction ratios of this line are its denominators

$$3k, 1, -5 = a_2, b_2, c_2$$

(\Rightarrow a vector along this line is $\vec{b}_2 = 3k\hat{i} + \hat{j} - 5\hat{k}$)

Because the lines are given to be perpendicular, therefore

$$\vec{b}_1 \cdot \vec{b}_2 (= a_1a_2 + b_1b_2 + c_1c_2) = 0$$

$$\Rightarrow (-3)(3k) + (2k)(1) + 2(-5) = 0$$

$$\Rightarrow -9k + 2k - 10 = 0$$

$$\Rightarrow -7k = 10 \Rightarrow k = \frac{-10}{7}$$

7. Find the vector equation of the line passing through (1, 2, 3)

and perpendicular to the plane $\vec{r} \cdot (\hat{i} + 2\hat{j} - 5\hat{k}) + 9 = 0$.


Sol. The required line passes through the point P(1, 2, 3).

\therefore Position vector \vec{a} (say) of point P is (1, 2, 3)

$$\Rightarrow \vec{a} = \hat{i} + 2\hat{j} + 3\hat{k}$$

Equation of the given plane is

$$\vec{r} \cdot (\hat{i} + 2\hat{j} - 5\hat{k}) + 9 = 0$$



$$\Rightarrow \vec{r} \cdot (\hat{i} + 2\hat{j} - 5\hat{k}) = -9 \quad r \cdot (i + 2j - 5k) + 9 = 0$$

Comparing with $\vec{r} \cdot \vec{n} = d$, we have normal vector \vec{n} to the given plane is $\vec{n} = \hat{i} + 2\hat{j} - 5\hat{k}$.

Because required line is perpendicular to the given plane,



therefore vector \vec{b} along the required line PM is $\vec{b} = \hat{i} + \hat{j} + \hat{k}$

$$2\hat{j} - 5\hat{k}$$

$$\therefore \text{Equation of required line is } \vec{r} = \vec{a} + \lambda \vec{b}$$

$$\text{i.e., } \vec{r} = \hat{i} + 2\hat{j} + 3\hat{k} + \lambda(\hat{i} + 2\hat{j} - 5\hat{k})$$

8. Find the equation of the plane passing through (a, b, c) and parallel to the plane $\vec{r} \cdot (\hat{i} + \hat{j} + \hat{k}) = 2$.

Sol. Equation of the given plane is $\vec{r} \cdot (\hat{i} + \hat{j} + \hat{k}) = 2$

$$\Rightarrow (x\hat{i} + y\hat{j} + z\hat{k}) \cdot (\hat{i} + \hat{j} + \hat{k}) = 2$$

$$\left[\therefore \vec{r} = x\hat{i} + y\hat{j} + z\hat{k} \right]$$

$$\Rightarrow x + y + z = 2$$

\therefore Equation of any plane parallel to this plane is $x + y + z = \lambda$ (i) (changing constant term only)

To find λ : Plane (i) passes through the point (a, b, c) (given). Putting $x = a, y = b, z = c$ in (i) $a + b + c = \lambda$

Putting $\lambda = a + b + c$ in (i), equation of required plane is $x + y + z = a + b + c$

9. Find the shortest distance between the lines

$$\vec{r} = 6\hat{i} + 2\hat{j} + 2\hat{k} + \lambda(\hat{i} - 2\hat{j} + 2\hat{k})$$

and $\vec{r} = -4\hat{i} - \hat{k} + \mu(3\hat{i} - 2\hat{j} - 2\hat{k})$

Sol. Given: vector equation of one line is

$$\vec{r} = 6\hat{i} + 2\hat{j} + 2\hat{k} + \lambda(\hat{i} - 2\hat{j} + 2\hat{k})$$

Comparing with $\vec{r} = \vec{a}_1 + \lambda \vec{b}_1$ we have



$$\vec{a}_1 = 6\hat{i} + 2\hat{j} + 2\hat{k} \quad \text{and} \quad \vec{b}_1 = \hat{i} - 2\hat{j} + 2\hat{k}$$

Given: Vector equation of second line is

$$\vec{r} = -4\hat{i} - \hat{k} + \mu(3\hat{i} - 2\hat{j} - 2\hat{k})$$



Comparing with $\vec{r} = \vec{a}_2 + \mu \vec{b}_2$, we have

$$\vec{a}_2 = -4\hat{i} - \hat{k} \quad \text{and} \quad \vec{b}_2 = 3\hat{i} - 2\hat{j} - 2\hat{k}$$

We know that length of shortest distance between two (skew) lines is

$$\frac{|\vec{(a}_2 - \vec{a}_1) \cdot (\vec{b}_1 \times \vec{b}_2)|}{|\vec{b}_1 \times \vec{b}_2|} \quad \dots(i)$$

$$\begin{aligned} \text{Now } \vec{a}_2 - \vec{a}_1 &= -4\hat{i} - \hat{k} - (6\hat{i} + 2\hat{j} + 2\hat{k}) \\ &= -4\hat{i} - \hat{k} - 6\hat{i} - 2\hat{j} - 2\hat{k} = -10\hat{i} - 2\hat{j} - 3\hat{k} \end{aligned}$$

$$\text{Again } \vec{b}_1 \times \vec{b}_2 = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & -2 & 2 \\ 3 & -2 & -2 \end{vmatrix}$$

Expanding along first row

$$= \hat{i}(4 + 4) - \hat{j}(-2 - 6) + \hat{k}(-2 + 6) = 8\hat{i} + 8\hat{j} + 4\hat{k}$$

$$\begin{aligned} \therefore (\vec{a}_2 - \vec{a}_1) \cdot (\vec{b}_1 \times \vec{b}_2) &= (-10)8 + (-2)8 + (-3)4 \\ &= -80 - 16 - 12 = -108 \end{aligned}$$

$$\begin{aligned} \text{and } |\vec{b}_1 \times \vec{b}_2| &= \sqrt{(8)^2 + (8)^2 + (4)^2} \\ &= \sqrt{64 + 64 + 16} = \sqrt{144} = 12 \end{aligned}$$

Putting these values in (i), length of shortest distance

$$= \frac{|-108|}{12} = \frac{108}{12} = 9.$$

10. Find the coordinates of the point where the line through (5, 1, 6) and (3, 4, 1) crosses the YZ-plane.

Sol. Given: A line through the points A(5, 1, 6) and B(3, 4, 1).

\therefore Direction ratios of this line AB are


$$x_2 - x_1, y_2 - y_1, z_2 - z_1$$

$$\text{i.e., } 3 - 5, 4 - 1, 1 - 6$$

$$\text{i.e., } -2, 3, -5 = a, b, c$$

\therefore Equation of line AB is

DS CUET Academy



$$\frac{z-z_1}{a} = \frac{y-y_1}{b} = \frac{x-x_1}{c}$$

YZ-plane $\Rightarrow x = 0$

$$\text{i.e., } \frac{z-5}{-2} = \frac{y-1}{3} = \frac{x-6}{-5} \quad \dots(i)$$



Let us find the coordinates of the point where this line AB crosses (\Rightarrow cuts or meets) the YZ-plane ($\Rightarrow x = 0$) ...*(ii)*
To find this point P, let us solve (i) and (ii) for x, y, z .

$$\text{Putting } x = 0 \text{ from (ii) in (i), } \frac{-5}{-2} = \frac{y-1}{3} = \frac{x-6}{-5}$$

$$\Rightarrow \frac{5}{2} = \frac{y-1}{3} = \frac{x-6}{-5}$$

$$\Rightarrow \frac{y-1}{3} = \frac{5}{2} \quad \text{and} \quad \frac{x-6}{-5} = \frac{5}{2}$$

$$\Rightarrow 2y - 2 = 15 \quad \text{and} \quad 2z - 12 = -25$$

$$\Rightarrow 2y = 17 \quad \text{and} \quad 2z = -13$$

$$\Rightarrow y = \frac{17}{2} \quad \text{and} \quad z = \frac{-13}{2}$$

$$\therefore \text{ Required point is } P\left(0, \frac{17}{2}, \frac{-13}{2}\right)$$

11. Find the coordinates of the point where the line through (5, 1, 6) and (3, 4, 1) crosses the ZX-plane.

Sol. Given: A line through the points A(5, 1, 6) and B(3, 4, 1).

For figure, see the figure for Q. No. 10

\therefore Direction ratios of this line AB are

$$\frac{x_2 - x_1}{a}, \frac{y_2 - y_1}{b}, \frac{z_2 - z_1}{c} \quad \text{i.e., } \frac{3 - 5}{-2}, \frac{4 - 1}{3}, \frac{1 - 6}{-5}$$

$$\therefore \text{ Equation of line AB is } \frac{z - z_1}{a} = \frac{y - y_1}{b} = \frac{x - x_1}{c}$$

$$\text{i.e., } \frac{z - 5}{-2} = \frac{y - 1}{3} = \frac{x - 6}{-5} \quad \dots(i)$$

Let us find the coordinates of the point where this line AB crosses (\Rightarrow cuts or meets) the ZX-plane ($\Rightarrow y = 0$) ...*(ii)*

To find this point P, let us solve (i) and (ii) for x, y, z .

Putting $y = 0$ from (ii) in (i), we have

$$\frac{z-5}{-2} = \frac{-1}{3} = \frac{x-6}{-5} \quad \Rightarrow \quad \frac{z-5}{-2} = \frac{-1}{3} \quad \text{and} \quad \frac{x-6}{-5} = \frac{-1}{3}$$

$$\Rightarrow 3x - 15 = 2 \quad \text{and} \quad 3z - 18 = 5$$

$$\Rightarrow 3x = 17 \quad \text{and} \quad 3z = 23$$

$$\Rightarrow x = \frac{17}{3} \quad \text{and} \quad z = \frac{23}{3}$$

\therefore Required point is $P\left(\frac{17}{3}, 0, \frac{23}{3}\right)$

12. Find the coordinates of the point where the line through $(3, -4, -5)$ and $(2, -3, 1)$ crosses the plane $2x + y + z = 7$.

Sol. Direction ratios of the line joining the points

$A(3, -4, -5)$ and $B(2, -3, 1)$ are
 $2 - 3, -3 - (-4), 1 - (-5)$ i.e., $-1, 1, 6$



$$\therefore \text{Equations of the line AB are } \frac{z-3}{-1} = \frac{y+4}{1} = \frac{x+5}{6} \quad \dots(i)$$

$$\left| \frac{z - z_1}{a} = \frac{y - y_1}{b} = \frac{x - x_1}{c} \right|$$

Equation of the plane is $2x + y + z = 7$... (ii)

Let us find the point where line (i) crosses (i.e., cuts i.e., meets) plane (ii).

\therefore For this, let us solve (i) and (ii) for x, y, z

$$\text{From (i), } \frac{z-3}{-1} = \frac{y+4}{1} = \frac{x+5}{6} = \lambda \text{ (say)}$$

$$\therefore \quad x - 3 = -\lambda, \quad y + 4 = \lambda, \quad z + 5 = 6\lambda$$

$$\therefore \quad x = 3 - \lambda, \quad y = -4 + \lambda, \quad z = -5 + 6\lambda \quad \dots(iii)$$

Putting these values of x, y, z in Eqn. (ii), we have

$$2(3 - \lambda) + (-4 + \lambda) + (-5 + 6\lambda) = 7$$

$$\text{or } 6 - 2\lambda - 4 + \lambda - 5 + 6\lambda = 7 \text{ or } 5\lambda = 10 \text{ or } \lambda = 2.$$

Putting $\lambda = 2$ in (iii), point of intersection of line (i) and plane (ii) is

$$x = 3 - 2 = 1, \quad y = -4 + 2 = -2, \quad z = -5 + 12 = 7.$$

\therefore Required point of intersection is $(1, -2, 7)$.

13. Find the equation of the plane passing through the point $(-1, 3, 2)$ and perpendicular to each of the planes $x + 2y + 3z = 5$ and $3x + 3y + z = 0$.

Sol. We know that equation of any plane through the point $(-1, 3, 2)$ is

$$a(x - x_1) + b(y - y_1) + c(z - z_1) = 0$$

$$\text{i.e., } a(x + 1) + b(y - 3) + c(z - 2) = 0 \quad \dots(ii)$$

$$\text{i.e., } ax + a + by - 3b + cz - 2c = 0$$

$$\text{or } ax + by + cz = -a + 3b + 2c$$

Given: This required plane is perpendicular to the plane

$$x + 2y + 3z = 5 \quad \therefore a_1a_2 + b_1b_2 + c_1c_2 = 0$$

i.e., Product of coefficients of $x + \dots = 0$

$$\therefore a(1) + b(2) + c(3) = 0 \quad \dots(iii)$$

Given: Again the required plane is perpendicular to the plane

$$3x + 3y + z = 0$$

$$\therefore a_1a_2 + b_1b_2 + c_1c_2 = 0$$

$$\text{i.e., } a(3) + b(3) + c(1) = 0 \quad \dots(iii)$$

Solving (ii) and (iii) for a, b, c

$$\frac{a}{2-9} = \frac{-b}{1-9} = \frac{c}{3-6} \quad \text{i.e., } \frac{a}{-7} = \frac{b}{8} = \frac{c}{-3}$$

Putting these value of a, b, c in (i), equation of required plane is

$$-7(x + 1) + 8(y - 3) + c(z - 2) = 0$$

$$\text{i.e.,} \quad -7x - 7 + 8y - 24 - 3z + 6 = 0$$

$$\text{i.e.,} \quad -7x + 8y - 3z - 25 = 0$$

$$\text{i.e.,} \quad 7x - 8y + 3z + 25 = 0.$$



14. If the points $(1, 1, p)$ and $(-3, 0, 1)$ be equidistant from the plane $\vec{r} \cdot (3\hat{i} + 4\hat{j} - 12\hat{k}) + 13 = 0$, then find the value of p .

Sol. Equation of the given plane is $\vec{r} \cdot (3\hat{i} + 4\hat{j} - 12\hat{k}) + 13 = 0$
 i.e., $(x\hat{i} + y\hat{j} + z\hat{k}) \cdot (3\hat{i} + 4\hat{j} - 12\hat{k}) + 13 = 0$
 $[\because \vec{r} = \text{Position vector of any point } (x, y, z)$
 on the plane $= x\hat{i} + y\hat{j} + z\hat{k}]$

$$\Rightarrow 3x + 4y - 12z + 13 = 0 \quad \dots(i)$$

Given: The points $(1, 1, p)$ and $(-3, 0, 1)$ are equidistant from plane (i).

\Rightarrow (Perpendicular) distance of point $(1, 1, p)$ from plane (i)
 = Distance of point $(-3, 0, 1)$ from plane (i)

$$\Rightarrow \frac{|3(1) + 4(1) - 12(p) + 13|}{\sqrt{9 + 16 + 144}} = \frac{|3(-3) + 4(0) - 12(1) + 13|}{\sqrt{9 + 16 + 144}}$$

$$\frac{|ax_1 + by_1 + cx_1 + d|}{\sqrt{a^2 + b^2 + c^2}}$$

$$\Rightarrow \frac{|3 + 4 - 12p + 13|}{13} = \frac{|-9 - 12 + 13|}{13}$$

$$\Rightarrow |20 - 12p| = |-8| = 8$$

$$\Rightarrow 20 - 12p = \pm 8 \quad [\because \text{If } |x| = a, a \geq 0, \text{ then } x = \pm a]$$

$$\text{Taking positive sign } 20 - 12p = 8 \Rightarrow -12p = -12$$

$$\Rightarrow p = 1$$

$$\text{Taking negative sign } 20 - 12p = -8$$

$$\Rightarrow -12p = -28 \Rightarrow p = \frac{-28}{-12} = \frac{7}{3} \text{ Hence, } p = 1 \text{ or } p = \frac{7}{3}$$

15. Find the equation of the plane passing through the line of intersection of the planes $\vec{r} \cdot (\hat{i} + \hat{j} + \hat{k}) = 1$ and

$\vec{r} \cdot (2\hat{i} + 3\hat{j} - \hat{k}) + 4 = 0$ and parallel to x-axis.


Sol. **Given:** Equation of first plane is

$$\vec{r} \cdot (\hat{i} + \hat{j} + \hat{k}) = 1$$

$$\Rightarrow (x\hat{i} + y\hat{j} + z\hat{k}) \cdot (\hat{i} + \hat{j} + \hat{k}) = 1$$

$$\Rightarrow x + y + z = 1$$

Making R.H.S. zero



$$\Rightarrow x + y + z - 1 = 0 \quad \dots(i)$$

Equation of second plane is



$$\begin{aligned} &\vec{r} \cdot (2\hat{i} + 3\hat{j} - \hat{k}) + 4 = 0 \\ \Rightarrow (x\hat{i} + y\hat{j} + z\hat{k}) \cdot (2\hat{i} + 3\hat{j} - \hat{k}) + 4 &= 0 \\ \Rightarrow 2x + 3y - z + 4 &= 0 \end{aligned} \quad \dots(ii)$$

(Here R.H.S. is already zero)

We know that equation of any plane passing through the line of intersection of these two planes is L.H.S. of (i) + λ [L.H.S. of (ii)] = 0

$$\begin{aligned} \text{i.e. } x + y + z - 1 + \lambda (2x + 3y - z + 4) &= 0 \\ \Rightarrow x + y + z - 1 + 2\lambda x + 3\lambda y - \lambda z + 4\lambda &= 0 \\ \Rightarrow (1 + 2\lambda)x + (1 + 3\lambda)y + (1 - \lambda)z - 1 + 4\lambda &= 0 \quad \dots(iii) \\ \Rightarrow ax + by + cz + d &= 0 \end{aligned}$$

Given : Required plane (iii) is parallel to x-axis $\Rightarrow a\hat{i} + b\hat{j} + c\hat{k} = 0$

$$\begin{aligned} \text{we know that a vector } \vec{b} \text{ along x-axis is } & b\hat{i} + 0\hat{j} + 0\hat{k} \\ \therefore \text{ D.R.'s of x-axis are } 1, 0, 0 \text{ | coeff of } \hat{i}, \hat{j}, \hat{k} &= l, m, n \end{aligned} \quad \dots(iv)$$

\therefore D.R.'s of x-axis are 1,0,0 | coeff of $\hat{i}, \hat{j}, \hat{k} = l, m, n$

Putting values of a,b,c, l,m,n, in (iv), we have

$$(1 + 2\lambda) \cdot 1 + (1 + 3\lambda) \cdot 0 + (1 - \lambda) \cdot 0 = 0$$

$$\Rightarrow 1 + 2\lambda = 0 \Rightarrow 2\lambda = -1 \Rightarrow \lambda = -\frac{1}{2}$$

Putting $\lambda = -\frac{1}{2}$ in (iii), Equation of required plane is

$$(1 - 1)x + \left(1 - \frac{3}{2}\right)y + \left(1 + \frac{1}{2}\right)z - 1 - 2 = 0$$

$$\text{or } -\frac{y}{2} + \frac{3z}{2} - 3 = 0$$

Multiplying by -2, $y - 3z + 6 = 0$

Note: Condition $a\hat{i} + b\hat{j} + c\hat{k} = 0$ is cartesian equivalent of

$$\vec{b} \cdot \vec{n} = 0$$

16. If O be the origin and the coordinates of P be (1, 2, -3), then find the equation of the plane passing through P and perpendicular to OP. $\times O(0, 0, 0)$

Sol. Given: Origin O(0, 0, 0) and point P(1, 2, -3).

We are to find the equation of the plane passing through P(1, 2, -3) = (x, y, z) and perpendicular to OP.

$$\times P(1, 2, -3)$$

∴ Direction ratios of normal OP to the plane are

$$1 - 0, 2 - 0, -3 - 0 \quad | \quad x_2 - x_1, y_2 - y_1, z_2 - z_1$$

i.e., $1, 2, -3 = a, b, c$.

∴ Equation of the required plane is



$$a(x - x_1) + b(y - y_1) + c(z - z_1) = 0$$

$$\text{i.e., } 1(x - 1) + 2(y - 2) - 3(z + 3) = 0$$

$$\text{i.e., } x - 1 + 2y - 4 - 3z - 9 = 0$$

$$\text{i.e., } x + 2y - 3z - 14 = 0.$$

17. Find the equation of the plane which contains the line of intersection of the planes $\vec{r} \cdot (\hat{i} + 2\hat{j} + 3\hat{k}) - 4 = 0$,
 $\vec{r} \cdot (2\hat{i} + \hat{j} - \hat{k}) + 5 = 0$ and which is perpendicular to the plane $\vec{r} \cdot (5\hat{i} + 3\hat{j} - 6\hat{k}) + 8 = 0$.

Sol. Equation of first plane is $\vec{r} \cdot (\hat{i} + 2\hat{j} + 3\hat{k}) - 4 = 0$

$$\text{i.e. } (x\hat{i} + y\hat{j} + z\hat{k}) \cdot (\hat{i} + 2\hat{j} + 3\hat{k}) - 4 = 0$$

$$\text{i.e. } x + 2y + 3z - 4 = 0 \quad \dots (i) \text{ (R.H.S. already zero)}$$

Equation of second plane is $\vec{r} \cdot (2\hat{i} + \hat{j} - \hat{k}) + 5 = 0$

$$\text{i.e. } (x\hat{i} + y\hat{j} + z\hat{k}) \cdot (2\hat{i} + \hat{j} - \hat{k}) + 5 = 0$$

$$\text{i.e. } 2x + y - z + 5 = 0 \quad \dots (ii) \text{ (R.H.S. already zero)}$$

We know that equation of any plane passing through the line of intersection of planes (i) and (ii) is

L.H.S. of (i) + λ L.H.S. of (ii) = 0

$$\text{i.e. } x + 2y + 3z - 4 + \lambda(2x + y - z + 5) = 0$$

$$\text{i.e. } x + 2y + 3z - 4 + 2\lambda x + \lambda y - \lambda z + 5\lambda = 0$$

$$\text{i.e. } (1 + 2\lambda)x + (2 + \lambda)y + (3 - \lambda)z - 4 + 5\lambda = 0 \quad \dots (iii)$$

Given : Required plane (iii) is perpendicular to the plane

$$\vec{r} \cdot (5\hat{i} + 3\hat{j} - 6\hat{k}) + 8 = 0$$

$$\text{i.e. } (x\hat{i} + y\hat{j} + z\hat{k}) \cdot (5\hat{i} + 3\hat{j} - 6\hat{k}) + 8 = 0$$

$$\text{i.e. } 5x + 3y - 6z + 8 = 0 \quad \dots (iv)$$

$$\therefore a_1a_2 + b_1b_2 + c_1c_2 = 0$$

$$\text{i.e. product of coefficient of } x \text{ in (iii) and (iv) } \dots + \dots = 0$$

$$\therefore (1 + 2\lambda)5 + (2 + \lambda)3 + (3 - \lambda)(-6) = 0$$

$$\Rightarrow 5 + 10\lambda + 6 + 3\lambda - 18 + 6\lambda = 0$$

$$\Rightarrow 19\lambda - 7 = 0 \Rightarrow 19\lambda = 7 \Rightarrow \lambda = \frac{7}{19}$$

Putting $\lambda = \frac{7}{19}$ in (iii), equation of required plane is

$$\left(1 + \frac{14}{19}\right)x + \left(2 + \frac{7}{19}\right)y + \left(3 - \frac{7}{19}\right)z - 4 + \frac{35}{19} = 0$$


Multiplying by 19, equation of required plane is

$$33x + 45y + 50z = 41$$

18. Find the distance of the point $(-1, -5, -10)$ from the point of intersection of the line

$$\vec{r} = 2\hat{i} - \hat{j} + 2\hat{k} + \lambda(3\hat{i} + 4\hat{j} + 2\hat{k}) \text{ and the plane } \vec{r} \cdot (\hat{i} - \hat{j} + \hat{k}) = 5.$$

Sol. Given: Point P (say) $(-1, -5, -10)$.



Given: Vector Equation of the line is

$$\vec{r} = 2\hat{i} - \hat{j} + 2\hat{k} + \lambda(3\hat{i} + 4\hat{j} + 2\hat{k}) = \vec{a} + \lambda \vec{b}$$

This line passes through the point $\vec{a} = 2\hat{i} - \hat{j} + 2\hat{k}$
 $= (2, -1, 2) = (x_1, y_1, z_1)$ and is parallel to the vector
 $3\hat{i} + 4\hat{j} + 2\hat{k}$ i.e. has d.r.'s 3,4,2.

\therefore Equation of the given line in cartesian form is

$$\frac{x-2}{3} = \frac{y+1}{4} = \frac{z-2}{2} = \lambda \text{ (say)} \quad \dots (i)$$

$$\left(\frac{x-x_1}{a} = \frac{y-y_1}{b} = \frac{z-z_1}{c} \right)$$

Vector equation of the given plane is $\vec{r} \cdot (\hat{i} - \hat{j} + \hat{k}) = 5$

$$\text{i.e. } (x\hat{i} + y\hat{j} + z\hat{k}) \cdot (\hat{i} - \hat{j} + \hat{k}) = 5$$

$$\Rightarrow x - y + z = 5 \quad \dots (ii)$$

Let us solve (i) and (ii) for x, y, z to find point of intersection (say Q) of line (i) and plane (ii).

From (i) $x - 2 = 3\lambda, y + 1 = 4\lambda, z - 2 = 2\lambda$

i.e. $x = 2 + 3\lambda, y = -1 + 4\lambda, z = 2 + 2\lambda$

\therefore Point Q is $(2 + 3\lambda, -1 + 4\lambda, 2 + 2\lambda)$ for some real λ ... (iii)

Putting these values of x, y, z in (ii)

$$2 + 3\lambda + 1 - 4\lambda + 2 + 2\lambda = 5 \Rightarrow \lambda = 0$$

Putting $\lambda = 0$ in (iii), point Q is $(2, -1, 2)$

\therefore Distance of the given point $P(-1, -5, -10)$ from the point $Q(2, -1, 2)$




$$\begin{aligned}
 &= PQ = \sqrt{(2+1)^2 + (-1+5)^2 + (2+10)^2} \\
 &\qquad\qquad\qquad (\sqrt{(z_2 - z_1)^2 + (y_2 - y_1)^2 + (x_2 - x_1)^2}) \\
 &= \sqrt{9+16+144} = \sqrt{169} = 13.
 \end{aligned}$$

19. Find the vector equation of the line passing through (1, 2, 3) and parallel to the planes

$$\vec{r} \cdot (\hat{i} - \hat{j} + 2\hat{k}) = 5 \text{ and } \vec{r} \cdot (3\hat{i} + \hat{j} + \hat{k}) = 6.$$

Sol. Given: The required line passes through the point A(1, 2, 3) (\vec{a})

$$\begin{aligned}
 \Rightarrow \vec{a} &= \text{Position vector of point A} \\
 &= \hat{i} + 2\hat{j} + 3\hat{k}
 \end{aligned}$$



Let \vec{b} be any vector along the required line.

$$\begin{aligned}
 \therefore \text{Vector equation of required line is } \vec{r} &= \vec{a} + \lambda \vec{b} \\
 \Rightarrow \vec{r} &= (\hat{i} + 2\hat{j} + 3\hat{k}) + \lambda \vec{b} \qquad \dots(i)
 \end{aligned}$$

Because the required line is parallel to the plane

$$\vec{r} \cdot (\hat{i} - \hat{j} + 2\hat{k}) = 5$$

$$(\text{Form } \vec{r} \cdot \vec{n}_1 = d_1 \text{ where } \vec{n}_1 = \hat{i} - \hat{j} + 2\hat{k}) \therefore \vec{b} \cdot \vec{n}_1 = 0$$

Similarly, $\vec{b} \cdot \vec{n}_2 = 0$

$$\begin{aligned}
 [\because \text{The required line is also parallel to the plane} \\
 \vec{r} \cdot (3\hat{i} + \hat{j} + \hat{k}) = 6 \quad \text{i.e., } \vec{r} \cdot \vec{n}_2 = d_2 \text{ where } \vec{n}_2 = 3\hat{i} + \hat{j} + \hat{k}]
 \end{aligned}$$

$$\text{Now } \vec{b} \cdot \vec{n}_1 = 0 \text{ and } \vec{b} \cdot \vec{n}_2 = 0$$

$\Rightarrow \vec{b}$ is perpendicular to both \vec{n}_1 and \vec{n}_2

$$\Rightarrow \vec{b} = \vec{n}_1 \times \vec{n}_2 \quad (\text{By definition of cross-product})$$

$$= \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & -1 & 2 \\ 3 & 1 & 1 \end{vmatrix}$$

Expanding along first row

$$\vec{b} = \hat{i}(-1-2) - \hat{j}(1-6) + \hat{k}(1+3) = -3\hat{i} + 5\hat{j} + 4\hat{k}$$



Putting this value of \vec{b} in (i), vector equation of required line is


$$\vec{r} = (\hat{i} + 2\hat{j} + 3\hat{k}) + \lambda(-3\hat{i} + 5\hat{j} + 4\hat{k}).$$

20. Find the vector equation of the line passing through the point $(1, 2, -4)$ and perpendicular to the two lines:

$$\frac{x-8}{3} = \frac{y+19}{-16} = \frac{z-10}{7} \quad \text{and} \quad \frac{x-15}{3} = \frac{y-29}{8} = \frac{z-5}{-5}.$$

Sol. Given: A point on the required line is $A(1, 2, -4)$.

$$\begin{aligned} \therefore \text{Position vector of point A is } \vec{a} &= (1, 2, -4) \\ &= \hat{i} + 2\hat{j} - 4\hat{k} \end{aligned}$$



Equations of the two given lines are

$$\frac{z-8}{3} = \frac{y+19}{-16} = \frac{x-10}{7} \quad (\text{Standard Form})$$

$$\text{and } \frac{z-15}{3} = \frac{y-29}{8} = \frac{x-5}{-5} \quad (\text{Standard Form})$$

\therefore Direction ratios of first given line are its denominators 3, -16, 7

i.e., a vector along this line is $\vec{b}_1 = 3\hat{i} - 16\hat{j} + 7\hat{k}$ and

direction ratios of the second given line are also its denominators 3, 8, -5

i.e., a vector along the second line is $\vec{b}_2 = 3\hat{i} + 8\hat{j} - 5\hat{k}$

Let \vec{b} be the vector along the required line perpendicular to the two given lines.

\therefore By definition of cross product

$$\vec{b} = \vec{b}_1 \times \vec{b}_2$$

$$\begin{array}{l} \rightarrow \quad \rightarrow \quad \rightarrow \\ \text{i.e., } \vec{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 3 & -16 & 7 \\ 3 & 8 & -5 \end{vmatrix} \end{array}$$

Expanding along first row

$$= \hat{i}(80 - 56) - \hat{j}(-15 - 21) + \hat{k}(24 + 48)$$



$$= 24 \hat{i} + 36 \hat{j} + 72 \hat{k}, \mathbf{b} = 12(2 \hat{i} + 3 \hat{j} + 6 \hat{k})$$

∴ Equation of the required line is $\mathbf{r} = \mathbf{a} + \lambda \mathbf{b}$

Putting values of \hat{i} and \hat{j} ,

$$\mathbf{r} = (\hat{i} + 2 \hat{j} - 4 \hat{k}) + \lambda(12)(2 \hat{i} + 3 \hat{j} + 6 \hat{k})$$

Replacing 12λ by λ ,

$$\mathbf{r} = (\hat{i} + 2 \hat{j} - 4 \hat{k}) + \lambda(2 \hat{i} + 3 \hat{j} + 6 \hat{k}).$$

21. Prove that if a plane has the intercepts a, b, c and is at a distance of p units from the origin, then

$$\frac{1}{a^2} + \frac{1}{b^2} + \frac{1}{c^2} = \frac{1}{p^2}.$$

Sol. We know that equation of a plane making intercepts a, b, c (on the axes) is $\frac{x}{a} + \frac{y}{b} + \frac{z}{c} = 1$

or $\frac{x}{a} + \frac{y}{b} + \frac{z}{c} - 1 = 0$... (i)

Perpendicular distance of the origin (0, 0, 0) from plane (i) = p (given)

$$\therefore \frac{\left| \frac{0}{a} + \frac{0}{b} + \frac{0}{c} - 1 \right|}{\sqrt{\frac{1}{a^2} + \frac{1}{b^2} + \frac{1}{c^2}}} = p \quad \left| \frac{1ax_1 + by_1 + cz_1 - d}{\sqrt{a^2 + b^2 + c^2}} \right| = \frac{dI}{\sqrt{a^2 + b^2 + c^2}}$$

Squaring both sides, $\frac{1}{\frac{1}{a^2} + \frac{1}{b^2} + \frac{1}{c^2}} = p^2$

Cross-multiplying, $p^2 \left(\frac{1}{a^2} + \frac{1}{b^2} + \frac{1}{c^2} \right) = 1$

Dividing by p^2 , $\frac{1}{a^2} + \frac{1}{b^2} + \frac{1}{c^2} = \frac{1}{p^2}.$

Choose the correct answer in Exercises Q. 22 and 23.

22. Distance between the two planes:

$2x + 3y + 4z = 4$ and $4x + 6y + 8z = 12$ is

- (A) 2 units
- (B) 4 units
- (C) 8 units



Sol. Given: Equation of one

plane is

units.

$$2x + 3y + 4z = 4$$

or $2x + 3y + 4z - 4 = 0$

...(i) (Making R.H.S. zero)

$$(ax + by + cz + d_1 = 0)$$

Equation of second plane is

$$4x + 6y + 8z = 12 \quad \text{or} \quad 4x + 6y + 8z - 12 = 0$$



The two planes are parallel as $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$ is satisfied.

$$\left(\because \frac{2}{4} = \frac{3}{6} = \frac{4}{8} \right)$$

Dividing every term of second equation by 2 to make coefficients of x, y, z equal in the equations of the two planes

$$\text{i.e., } \begin{aligned} 2x + 3y + 4z - 6 &= 0 \\ (ax + by + cz + d_2 &= 0) \end{aligned} \quad \dots(ii)$$

We know that distance between parallel planes (i) and (ii)

$$= \frac{I d_1 - d_2 I}{\sqrt{a^2 + b^2 + c^2}} = \frac{I - 4 - (-6)I}{\sqrt{(2)^2 + (3)^2 + (4)^2}} = \frac{I - 4 + 6I}{\sqrt{4 + 9 + 16}} = \frac{2}{\sqrt{29}}$$

\therefore Option (D) is the correct answer.

23. The planes: $2x - y + 4z = 5$ and $5x - 2.5y + 10z = 6$ are

- (A) Perpendicular (B) Parallel
(C) intersect y -axis (D) passes through $\left(0, 0, \frac{5}{4} \right)$.

Sol. Equations of the given planes are

$$\begin{aligned} 2x - y + 4z &= 5 \\ (a_1x + b_1y + c_1z + d_1 &= 0) \\ 5x - 2.5y + 10z &= 6 \\ (a_2x + b_2y + c_2z + d_2 &= 0) \end{aligned}$$

Let us test option (A). Are these planes perpendicular?

$$\begin{aligned} a_1a_2 + b_1b_2 + c_1c_2 &= 2(5) + (-1)(-2.5) + 4(10) \\ &= 10 + 2.5 + 40 = 52.5 \neq 0 \end{aligned}$$

\therefore Planes are not perpendicular.

\therefore Option (A) is not correct answer.

Let us test option (B). Are these planes parallel?

$$\text{Here, } \frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$$

$$\Rightarrow \frac{2}{5} = \frac{-1}{-2.5} = \frac{4}{10}$$

$$\Rightarrow \frac{2}{5} = \frac{1}{\left(\frac{2.5}{10} \right)} = \frac{2}{5} \Rightarrow \frac{2}{5} = \frac{10}{25} = \frac{2}{5}$$

$$\Rightarrow \frac{2}{5} = \frac{2}{5} = \frac{2}{5} \text{ which is true. } \therefore \text{ Planes are parallel.}$$

\therefore Option (B) is correct answer.